

UCD GEARY INSTITUTE DISCUSSION PAPER SERIES

Analyzing Social Experiments as Implemented: A Reexamination of

the Evidence From the HighScope Perry Preschool Program

James Heckman, Seong Hyeok Moon, Rodrigo Pinto,

Peter Savelyev, and Adam Yavitz¹

University of Chicago

July 22, 2010

¹James Heckman is Henry Schultz Distinguished Service Professor of Economics at the University of Chicago, Professor of Science and Society, University College Dublin, Alfred Cowles Distinguished Visiting Professor, Cowles Foundation, Yale University, and Senior Fellow, American Bar Foundation. Seong Hyeok Moon, Rodrigo Pinto, Peter Savelyev and Adam Yavitz are graduate students at the University of Chicago. A version of this paper was presented at a seminar at the HighScope Perry Foundation, Ypsilanti, Michigan, December 2006; at a conference at the Minneapolis Federal Reserve in December 2007; at a conference on the role of early life conditions at the Michigan Poverty Research Center, University of Michigan, December 2007; at aJacobs Foundation conference at Castle Marbach, April 2008; at the Leibniz Network Conference on Noncognitive Skills in Mannheim, Germany, May 2008; at an Institute for Research on Poverty conference, Madison, Wisconsin, June 2008; and at a conference on early childhood at the Brazilian National Academy of Sciences, Rio de Janeiro, Brazil, December 2009. We thank the editor and two anonymous referees for helpful comments which greatly improved this draft of the paper. We have benefited from comments received on early drafts of this paper at two brown bag lunches at the Statistics Department, University of Chicago, hosted by Stephen Stigler. We thank all of the workshop participants. In addition, we thank Amanda Agan, Mathilde Almlund, Joseph Altonji, Ricardo Barros, Dan Black, Steve Durlauf, Chris Hansman, Tim Kautz, Paul LaFontaine, Devesh Raval, Azeem Shaikh, Je_ Smith, and Steve Stigler for helpful comments. Our collaboration with Azeem Shaikh on related work has greatly strengthened the analysis in this paper. This research was supported in part by the American Bar Foundation, the Committee for Economic Development; by a grant from the Pew Charitable Trusts and the Partnership for America's Economic Success; the JB & MK Pritzker Family Foundation; Susan Thompson Bu_ett Foundation; Mr. Robert Dugger; and NICHD R01HD043411. The views expressed in this presentation are those of the authors and not necessarily those of the funders listed here. Supplementary materials for this paper may be found at http://jenni.uchicago.edu/Perry/.



UCD GEARY INSTITUTE DISCUSSION PAPER SERIES

Analyzing Social Experiments as Implemented: A Reexamination of

the Evidence From the HighScope Perry Preschool Program

James Heckman, Seong Hyeok Moon, Rodrigo Pinto,

Peter Savelyev, and Adam Yavitz¹

University of Chicago

July 22, 2010

¹James Heckman is Henry Schultz Distinguished Service Professor of Economics at the University of Chicago, Professor of Science and Society, University College Dublin, Alfred Cowles Distinguished Visiting Professor, Cowles Foundation, Yale University, and Senior Fellow, American Bar Foundation. Seong Hyeok Moon, Rodrigo Pinto, Peter Savelyev and Adam Yavitz are graduate students at the University of Chicago. A version of this paper was presented at a seminar at the HighScope Perry Foundation, Ypsilanti, Michigan, December 2006; at a conference at the Minneapolis Federal Reserve in December 2007; at a conference on the role of early life conditions at the Michigan Poverty Research Center, University of Michigan, December 2007; at aJacobs Foundation conference at Castle Marbach, April 2008; at the Leibniz Network Conference on Noncognitive Skills in Mannheim, Germany, May 2008; at an Institute for Research on Poverty conference, Madison, Wisconsin, June 2008; and at a conference on early childhood at the Brazilian National Academy of Sciences, Rio de Janeiro, Brazil, December 2009. We thank the editor and two anonymous referees for helpful comments which greatly improved this draft of the paper. We have benefited from comments received on early drafts of this paper at two brown bag lunches at the Statistics Department, University of Chicago, hosted by Stephen Stigler. We thank all of the workshop participants. In addition, we thank Amanda Agan, Mathilde Almlund, Joseph Altonji, Ricardo Barros, Dan Black, Steve Durlauf, Chris Hansman, Tim Kautz, Paul LaFontaine, Devesh Raval, Azeem Shaikh, Je_ Smith, and Steve Stigler for helpful comments. Our collaboration with Azeem Shaikh on related work has greatly strengthened the analysis in this paper. This research was supported in part by the American Bar Foundation, the Committee for Economic Development; by a grant from the Pew Charitable Trusts and the Partnership for America's Economic Success; the JB & MK Pritzker Family Foundation; Susan Thompson Bu_ett Foundation; Mr. Robert Dugger; and NICHD R01HD043411. The views expressed in this presentation are those of the authors and not necessarily those of the funders listed here. Supplementary materials for this paper may be found at http://jenni.uchicago.edu/Perry/.

Analyzing Social Experiments as Implemented: A Reexamination of the

Evidence From the HighScope Perry Preschool Program

James Heckman, Seong Hyeok Moon, Rodrigo Pinto,

Peter Savelyev, and Adam Yavitz¹

University of Chicago

July 22, 2010

¹James Heckman is Henry Schultz Distinguished Service Professor of Economics at the University of Chicago, Professor of Science and Society, University College Dublin, Alfred Cowles Distinguished Visiting Professor, Cowles Foundation, Yale University, and Senior Fellow, American Bar Foundation. Seong Hyeok Moon, Rodrigo Pinto, Peter Savelyev and Adam Yavitz are graduate students at the University of Chicago. A version of this paper was presented at a seminar at the HighScope Perry Foundation, Ypsilanti, Michigan, December 2006; at a conference at the Minneapolis Federal Reserve in December 2007; at a conference on the role of early life conditions at the Michigan Poverty Research Center, University of Michigan, December 2007; at a Jacobs Foundation conference at Castle Marbach, April 2008; at the Leibniz Network Conference on Noncognitive Skills in Mannheim, Germany, May 2008; at an Institute for Research on Poverty conference, Madison, Wisconsin, June 2008; and at a conference on early childhood at the Brazilian National Academy of Sciences, Rio de Janeiro, Brazil, December 2009. We thank the editor and two anonymous referees for helpful comments which greatly improved this draft of the paper. We have benefited from comments received on early drafts of this paper at two brown bag lunches at the Statistics Department, University of Chicago, hosted by Stephen Stigler. We thank all of the workshop participants. In addition, we thank Amanda Agan, Mathilde Almlund, Joseph Altonji, Ricardo Barros, Dan Black, Steve Durlauf, Chris Hansman, Tim Kautz, Paul LaFontaine, Devesh Raval, Azeem Shaikh, Jeff Smith, and Steve Stigler for helpful comments. Our collaboration with Azeem Shaikh on related work has greatly strengthened the analysis in this paper. This research was supported in part by the American Bar Foundation, the Committee for Economic Development; by a grant from the Pew Charitable Trusts and the Partnership for America's Economic Success; the JB & MK Pritzker Family Foundation; Susan Thompson Buffett Foundation; Mr. Robert Dugger; and NICHD R01HD043411. The views expressed in this presentation are those of the authors and not necessarily those of the funders listed here. Supplementary materials for this paper may be found at http://jenni.uchicago.edu/Perry/.

Abstract

Social experiments are powerful sources of information about the effectiveness of interventions. In practice, initial randomization plans are almost always compromised. Multiple hypotheses are frequently tested. "Significant" effects are often reported with *p*-values that do not account for preliminary screening from a large candidate pool of possible effects. This paper develops tools for analyzing data from experiments as they are actually implemented.

We apply these tools to analyze the influential HighScope Perry Preschool Program. The Perry program was a social experiment that provided preschool education and home visits to disadvantaged children during their preschool years. It was evaluated by the method of random assignment. Both treatments and controls have been followed from age 3 through age 40.

Previous analyses of the Perry data assume that the planned randomization protocol was implemented. In fact, as in many social experiments, the intended randomization protocol was compromised. Accounting for compromised randomization, multiple-hypothesis testing, and small sample sizes, we find statistically significant and economically important program effects for both males and females. We also examine the representativeness of the Perry study.

Keywords: early childhood intervention; compromised randomization; social experiment; multiple-hypothesis testing.

JEL codes: I21, C93, J15, V16.

1 Introduction

Social experiments can produce valuable information about the effectiveness of interventions. However, many social experiments are compromised by departures from initial randomization plans.¹ Many have small sample sizes. Applications of large sample statistical procedures may produce misleading inferences. In addition, most social experiments have multiple outcomes. This creates the danger of selective reporting of "significant" effects from a large pool of possible effects, biasing downward reported p-values. This paper develops tools for analyzing the evidence from experiments with multiple outcomes as they are implemented rather than as they are planned. We apply these tools to reanalyze an influential social experiment.

The HighScope Perry Preschool program, conducted in the 1960s, was an early childhood intervention that provided preschool education to low-IQ, disadvantaged African-American children living in Ypsilanti, Michigan. The study was evaluated by the method of random assignment. Participants were followed through age 40 and plans are under way for an age-50 followup. The beneficial long-term effects reported for the Perry program constitute a cornerstone of the argument for early childhood intervention efforts throughout the world.

Many analysts discount the reliability of the Perry study. For example, Hanushek and Lindseth (2009), among others, claim that the sample size of the study is too small to make valid inferences about the program. Herrnstein and Murray (1994) claim that estimated effects of the program are small and that many are not statistically significant. Others express the concern that previous analyses selectively report statistically significant estimates, biasing the inference about the program (Anderson, 2008).

There is a potentially more devastating critique. As happens in many social experiments, the proposed randomization protocol for the Perry study was compromised. This compromise casts doubt on the validity of evaluation methods that do not account for the compromised randomization and calls into question the validity of the simple statistical

¹See the discussion in Heckman (1992); Hotz (1992); and Heckman, LaLonde, and Smith (1999).

procedures previously applied to analyze the Perry study.²

In addition, there is the question of how representative the Perry population is of the general African-American population. Those who advocate access to universal early child-hood programs often appeal to the evidence from the Perry study, even though the project only targeted a disadvantaged segment of the population.³

This paper develops and applies small-sample permutation procedures that are tailored to test hypotheses on samples generated from the less-than-ideal randomizations conducted in many social experiments. We apply these tools to the data from the Perry experiment. We correct estimated treatment effects for imbalances that arose in implementing the randomization protocol and from post-randomization reassignment. We address the potential problem that arises from arbitrarily selecting "significant" hypotheses from a set of possible hypotheses using recently developed stepdown multiple-hypothesis testing procedures. The procedures we use minimize the probability of falsely rejecting any true null hypotheses.

Using these tools, this paper demonstrates the following points: (a) Statistically significant Perry treatment effects survive analyses that account for the small sample size of the study. (b) Correcting for the effect of selectively reporting statistically significant responses, there are substantial impacts of the program on males and females. Results are stronger for females at younger adult ages and for males at older adult ages. (c) Accounting for the compromised randomization of the program strengthens the evidence for important program effects compared to the evidence reported in the previous literature that neglects the imbalances created by compromised randomization. (d) Perry participants are representative of a low-ability, disadvantaged African-American population.

This paper proceeds as follows. Section 2 describes the Perry experiment. Section 3

²This problem is pervasive in the literature. For example, in the Abecedarian program, randomization was also compromised as some initially enrolled in the experiment were later dropped (Campbell and Ramey, 1994). In the SIME-DIME experiment, the randomization protocol was never clearly described. See Kurz and Spiegelman (1972). Heckman, LaLonde, and Smith (1999) chronicle the variety of "threats to validity" encountered in many social experiments.

 $^{^{3}}$ See, for example, The Pew Center on the States (2009) for one statement about the benefits of universal programs.

discusses the statistical challenges confronted in analyzing the Perry experiment. Section 4 presents our methodology. Our main empirical analysis is presented in Section 5. Section 6 examines the representativeness of the Perry sample. Section 7 compares our analysis to previous analyses of Perry. Section 8 concludes. Supplementary material is placed in the Web Appendix.⁴

2 Perry: Experimental Design and Background

The HighScope Perry Program was conducted during the early- to mid-1960s in the district of the Perry Elementary School, a public school in Ypsilanti, Michigan, a town near Detroit. The sample size was small: 123 children allocated over five entry cohorts. Data were collected at age 3, the entry age, and through annual surveys until age 15, with additional follow-ups conducted at ages 19, 27, and 40. Program attrition remained low through age 40, with over 91% of the original subjects interviewed. Two-thirds of the attrited were dead. The rest were missing.⁵ Numerous measures were collected on economic, criminal, and educational outcomes over this span as well as on cognition and personality. Program intensity was low compared to that in many subsequent early childhood development programs.⁶ Beginning at age 3, and lasting 2 years, treatment consisted of a 2.5-hour educational preschool on weekdays during the school year, supplemented by weekly home visits by teachers.⁷ HighScope's innovative curriculum, developed over the course of the Perry experiment, was based on the principle of active learning, guiding students through the formation of key developmental factors using intensive child-teacher interactions (Schweinhart et al. 1993, pp. 34–36; Weikart et al. 1978, pp. 5–6, 21–23). A more complete description of the Perry program curriculum is given in Web Appendix A.⁸

⁴http://jenni.uchicago.edu/Perry/

⁵There are two missing controls and two missing treatments. Five controls and two treatments are dead. ⁶The Abecedarian program is an example (see, e.g., Campbell et al., 2002). Cunha, Heckman, Lochner, and Masterov (2006) and Reynolds and Temple (2008) discuss a variety of these programs and compare their intensity.

⁷An exception is that the first entry cohort received only 1 year of treatment, beginning at age 4.

⁸The website can be accessed at http://jenni.uchicago.edu/Perry/.

Eligibility Criteria The program admitted five entry cohorts in the early 1960s, drawn from the population surrounding the Perry Elementary school. Candidate families for the study were identified from a survey of the families of the students attending the elementary school, by neighborhood group referrals, and through door-to-door canvassing. The eligibility rules for participation were that the participants should (i) be African-American; (ii) have a low IQ (between 70 and 85) at study entry,⁹ and (iii) be disadvantaged as measured by parental employment level, parental education, and housing density (persons per room). The Perry study targeted families who were more disadvantaged than most other African-American families in the United States. but were representative of a large segment of the disadvantaged African-American population. We discuss the issue of the representativeness of the program compared to the general African-American population in Section 6.

Among children in the Perry Elementary School neighborhood, Perry study families were particularly disadvantaged. Table 1 shows that compared to other families with children in the Perry School catchment area, Perry study families were younger, had lower levels of parental education, and had fewer working mothers. Further, Perry program families had fewer educational resources, larger families, and greater participation in welfare, compared to the families with children in another neighborhood elementary school in Ypsilanti, the Erickson school, situated in a predominantly middle-class white neighborhood.

We do not know whether, among eligible families in the Perry catchment, those who volunteered to participate in the program were more motivated than other families, and whether this greater motivation would have translated into better child outcomes. However, according to Weikart, Bond, and McNeil (1978, p. 16), "virtually all eligible children were enrolled in the project," so this potential concern appears to be unimportant.

Randomization Protocol The randomization protocol used in the Perry study was complex. According to Weikart et al. (1978, p. 16), for each designated eligible entry cohort,

⁹Measured by the Stanford-Binet IQ test (1960s norming). The average IQ in the general population is 100 by construction. IQ range for Perry participants is 1-2 standard deviations below the average.

Table 1: Comparing Families of Participants with Other Families with Children i	n the
Perry Elementary School Catchment and a Nearby School in Ypsilanti, Michigan	

		Perry School (Overall) ^a	Perry Preschool ^b	Erickson School ^c
	Average Age	35	31	32
Mother	Mean Years of Education	10.1	9.2	12.4
	% Working	60%	$\frac{3.2}{20\%}$	12.4 15%
	Mean Occupational Level ^d	1.4	1.0	2.8
	% Born in South	77%	80%	2.0 22%
	% Educated in South	53%	48%	17%
	% Fathers Living in the Home	63%	48%	100%
Father	Mean Age	40	35	35
fat	Mean Years of Education	9.4	8.3	13.4
	Mean Occupational Level ^d	1.6	1.1	3.3
	Mean SES ^e	11.5	4.2	16.4
	Mean # of Children	3.9	4.5	3.1
	Mean $\#$ of Rooms	5.9	4.8	6.9
ne	Mean # of Others in Home	0.4	0.3	0.1
on	% on Welfare	30%	58%	0%
H	% Home Ownership	33%	5%	85%
S	% Car Ownership	64%	39%	98%
ily	% Members of Library ^f	25%	10%	35%
Family & Home	% with Dictionary in Home	65%	24%	91%
Ĩ	% with Magazines in Home	51%	43%	86%
	% with Major Health Problems	16%	13%	9%
	% Who Had Visited a Museum	20%	2%	42%
	% Who Had Visited a Zoo	49%	26%	72%
	N	277	45	148

Source: Weikart, Bond, and McNeil (1978). Notes: (a) These are data on parents who attended parentteacher meetings at the Perry school or who were tracked down at their homes by Perry personnel (Weikart, Bond, and McNeil, 1978, pp. 12–15); (b) The Perry Preschool subsample consists of the full sample (treatment and control) from the first two waves; (c) The Erickson School was an "all-white school located in a middleclass residential section of the Ypsilanti public school district." (ibid., p. 14); (d) Occupation level: 1 =unskilled; 2 = semiskilled; 3 = skilled; 4 = professional; (e) See the notes at the base of Figure 3 for the definition of socio-economic status (SES) index; (f) Any member of the family. children were assigned to treatment and control groups in the following way, which is graphically illustrated in Figure 1:

- In any entering cohort, younger siblings of previously enrolled families were assigned the same treatment status as their older siblings.¹⁰
- 2. Those remaining were ranked by their entry IQ scores.¹¹ Odd- and even-ranked subjects were assigned to two separate unlabeled groups.

Balancing on IQ produced an imbalance on family background measures. This was corrected in a second, "balancing", stage of the protocol.

- 3. Some individuals initially assigned to one group were swapped between the unlabeled groups to balance gender and mean socio-economic (SES) status, "with Stanford-Binet scores held more or less constant."
- 4. A flip of a coin (a single toss) labeled one group as "treatment" and the other as "control."
- 5. Some individuals provisionally assigned to treatment, whose mothers were employed at the time of the assignment, were swapped with control individuals whose mothers were not employed. The rationale for these swaps was that it was difficult for working mothers to participate in home visits assigned to the treatment group and because of transportation difficulties.¹² A total of five children of working mothers initially assigned to treatment were reassigned to control.

¹⁰The rationale for excluding younger siblings from the randomization process was that enrolling children in the same family in different treatment groups would weaken the observed treatment effect due to withinfamily spillovers.

¹¹Ties were broken by a toss of a coin.

¹²The following quotation from an early monograph on Perry summarizes the logic of the study planners: "Occasional exchanges of children between groups also had to be made because of the inconvenience of half-day preschool for working mothers and the transportation difficulties of some families. No funds were available for transportation or full-day care, and special arrangements could not always be made." (Weikart, Bond, and McNeil, 1978, p. 17)

Even after the swaps at stage 3 were made, pre-program measures were still somewhat imbalanced between treatment and control groups. See Figure 2 for IQ and Figure 3 for SES index.

3 Statistical Challenges in Analyzing the Perry Program

Drawing valid inference from the Perry study requires meeting three statistical challenges: (i) small sample size; (ii) compromise in the randomization protocol; and (iii) the large number of outcomes and associated hypotheses, which creates the danger of selectively reporting "significant" estimates out of a large candidate pool of estimates, thereby biasing downward reported p-values.

Small Sample Size The small sample size of the Perry study and the non-normality of many outcome measures call into question the validity of classical tests, such as those based on the t-, F-, and χ^2 -statistics.¹³ Classical statistical tests rely on central limit theorems and produce inferences based on p-values that are only asymptotically valid.


A substantial literature demonstrates that classical testing procedures can be unreliable when sample sizes are small and the data are non-normal.¹⁴ Both features characterize the Perry study. There are approximately 25 observations per gender in each treatment assignment group and the distribution of observed measures is often highly skewed.¹⁵ Our paper addresses the problem of small sample size by using permutation-based inference procedures that are valid in small samples.

The Treatment Assignment Protocol The randomization protocol implemented in the Perry study diverged from the original design. Treatment and control statuses were

 $^{^{13}\}mathrm{Heckman}$ (2005) raises this concern in the context of the Perry program.

¹⁴See Micceri (1989) for a survey.

¹⁵Crime measures are a case in point.



*This figure is a visual representation of the Perry Randomization Protocol. T and C refer to treatment and control groups respectively. Shaded circles represent males. Light circles represent females. G₁ and G₂ are unlabeled groups of participants.

Figure 1: Perry Randomization Protocol^{*}






Figure 2: IQ at Entry by Entry Cohort and Treatment Status

Note: Stanford-Binet IQ at study entry (age 3) was used to measure the baseline IQ.

reassigned for a subset of persons after an initial random assignment. This creates two potential problems.

First, such reassignments can induce correlation between treatment assignment and baseline characteristics of participants. If the baseline measures affect outcomes, treatment assignment can become correlated with outcomes through an induced common dependence. Such a relationship between outcomes and treatment assignment violates the assumption of independence between treatment assignment and outcomes in the absence of treatment effects. Moreover, reassignment produces an imbalance in the covariates between the treated and the controlled, as documented in Figures 2 and 3. For example, the working status of the mother was one basis for reassignment to the control group. Weikart, Bond, and McNeil (1978, p. 18) note that at baseline, children of working mothers had higher test scores. Not controlling for mother's working status would bias downward estimated treatment effects for schooling and other ability-dependent outcomes. We control for imbalances by conditioning on such covariates.





Notes: The socio-economic status (SES) index is a weighted linear combination of three variables: (a) average highest grade completed by whichever parent(s) was present, with a coefficient 0.5; (b) father's employment status (or mother's, if the father was absent): 3 for skilled, 2 for semi-skilled, and 1 for unskilled or none, all with a coefficient 2; (c) number of rooms in the house divided by number of people living in the household, with a coefficient 2. The skill level of the parent's job is rated by the study coordinators and is not clearly defined. An SES index of 11 or lower was the intended requirement for entry into the study (Weikart, Bond, and McNeil, 1978, p. 14). This criterion was not always adhered to: out of the full sample, 7 individuals had an SES index above the cutoff (6 out of 7 were in the treatment group, and 6 out of 7 were in the last two waves).

Second, even if treatment assignment is statistically independent of the baseline variables, compromised randomization can still produce biased inference. A compromised randomization protocol can generate treatment assignment distributions that differ from those that would result from implementation of the intended randomization protocol. As a consequence, incorrect inference can occur if the data are analyzed under the assumption that no compromise in randomization has occurred.

More specifically, analyzing the Perry study under the assumption that a fair coin decides the treatment assignment of each participant — as if an idealized, non-compromised randomization had occurred — mischaracterizes the actual treatment assignment mechanism and hence the probability of assignment to treatment. This can produce incorrect critical values and improper control of Type-I error. Section 4.5 presents a procedure that accounts for the compromised randomization using permutation-based inference conditioned on baseline background measures.

Multiple Hypotheses There are numerous outcomes reported in the Perry experiment. One has to be careful in conducting analyses to avoid selective reporting of statistically significant outcomes, as determined by single-hypothesis tests, without correcting for the effects of such preliminary screening on actual *p*-values. This practice is sometimes termed cherry picking.

Multiple-hypothesis testing procedures avoid bias in inference arising from selectively reporting statistically significant results by adjusting inference to take into account the overall set of outcomes from which the "significant" results are drawn.

The traditional approach to testing based on overall F-statistics involves testing the null hypothesis that *any* element of a block of hypotheses is rejected. We test that hypothesis as part of a general stepdown procedure, which also tests *which* hypotheses within the block of hypotheses are rejected.

Simple calculations suggest that concerns about the overall statistical significance of

	All Data	Male Subsample	Female Subsample
Percentage of p -values smaller than 1%	7%	3%	7%
Percentage of p -values smaller than 5%	23%	13%	22%
Percentage of $p\text{-values}$ smaller than 10%	34%	21%	31%

 Table 2: Percentage of Test Statistics Exceeding Various Significance Levels^a

^aBased on 715 outcomes in the Perry study. (See Schweinhart, Montie, Xiang, Barnett, Belfield, and Nores, 2005, for a description of the data.) 269 outcomes are from the period before the age-19 interview; 269 are from the age-19 interview; 95 are outcomes from the age-27 interview; 55 are outcomes from the age-40 interview.

treatment effects for the Perry study may have been overstated. Table 2 summarizes the inference for 715 Perry study outcomes by reporting the percentage of hypotheses rejected at various significance levels.¹⁶ If outcomes were statistically independent and there was no experimental treatment effect, we would expect only 1% of the hypotheses to be rejected at the 1% level, but instead 7% are rejected overall (3% for males and 7% for females). At the 5% significance level, we obtain a 23% overall rejection rate (13% for males and 22% for females). Far more than 10% of the hypotheses are statistically significant when the 10% level is used. These results suggest that treatment effects are present for each gender and for the full sample.

However, the assumption of independence among the outcomes used to make these calculations is quite strong. In our analysis, we use modern methods for testing multiple hypotheses that account for possible dependencies among outcomes. We use a stepdown multiple-hypothesis testing procedure that controls for the Family-Wise Error Rate (FWER) — the probability of rejecting at least one true null hypothesis among a set of hypotheses we seek to test jointly. This procedure is discussed below in Section 4.6.

 $^{^{16}}$ Inference is based on a permutation testing method where the *t*-statistic of the difference in means between treatment and control groups is used as the test statistic.

4 Methods

This section presents a framework for inference that addresses the problems raised in Section 3, namely, small samples, compromised randomization, and cherry picking. We first establish notation, discuss the benefits of a valid randomization, and consider the consequences of compromised randomization. We then introduce a general framework for representing randomized experiments. Using this framework, we develop a statistical framework for characterizing the conditions under which permutation-based inference produces valid small-sample inference when there is corruption of the intended randomization protocol. Finally, we discuss the multiple-hypothesis testing procedure used in this paper.

4.1 Randomized Experiments

The standard model of program evaluation describes the observed outcome for participant $i, Y_i, by Y_i = D_i Y_{i,1} + (1 - D_i) Y_{i,0}$, where $(Y_{i,0}, Y_{i,1})$ are potential outcomes corresponding to control and treatment status for participant i, respectively, and D_i is the assignment indicator: $D_i = 1$ if treatment occurs, $D_i = 0$ otherwise.

An evaluation problem arises because either $Y_{i,0}$ or $Y_{i,1}$ is observed, but not both. Selection bias can arise from participant self-selection into treatment and control groups so that sampled distributions of $Y_{i,0}$ and $Y_{i,1}$ are biased estimators of the population distributions. Properly implemented randomized experiments eliminate selection bias because they produce independence between $(Y_{i,0}, Y_{i,1})$ and D_i .¹⁷ Notationally, $(Y_0, Y_1) \perp D$, where Y_0, Y_1 , and D are vectors of variables across participants, and \perp denotes independence.

Selection bias can arise when experimenters fail to generate treatment groups that are comparable on unobserved background variables that affect outcomes. A properly conducted randomization avoids the problem of selection bias by inducing independence between unobserved variables and treatment assignments.

Compromised randomization can invalidate the assumption that $(Y_0, Y_1) \perp D$. The

 $^{^{17}\}mathrm{Web}$ Appendix B discusses this point in greater detail.

treatments and controls can have imbalanced covariate distributions.¹⁸ The following notational framework helps to clarify the basis for inference under compromised randomization that characterizes the Perry study.

4.2 Setup and Notation

Denote the set of participants by $\mathcal{I} = \{1, \ldots, I\}$, where I = 123 is the total number of Perry study participants. We denote the random vector representing treatment assignments by $D = (D_i; i \in \mathcal{I})$. The set \mathcal{D} is the support of the vector of random assignments, namely $\mathcal{D} = [0, 1] \times \cdots \times [0, 1]$, 123 times, so $\mathcal{D} = [0, 1]^{123}$. Define the pre-program variables used in the randomization protocol by $X = (X_i; i \in \mathcal{I})$. For the Perry study, baseline variables X consist of data on the following measures: IQ, enrollment cohort, socio-economic status (SES) index, family structure, gender, and maternal employment status, all measured at study entry.

Assignment to treatment is characterized by a function M. The arguments of M are variables that affect treatment assignment. Define R as a random vector that describes the outcome of a randomization device (e.g., a flip of a coin to assign treatment status). Prior to determining the realization of R, two groups are formed on the basis of pre-program variables X. Then R is realized and its value is used to assign treatment status. R does not depend on the composition of the two groups. After the initial treatment assignment, individuals are swapped across assigned treatment groups based on some observed background characteristics X (e.g., mother's working status). M captures all three aspects of the treatment assignment mechanism. The following assumptions formalize the treatment

¹⁸Heckman and Smith (1995), Heckman, LaLonde, and Smith (1999), and Heckman and Vytlacil (2007) discuss randomization bias and substitution bias. The Perry study does not appear to be subject to these biases. Randomization bias occurs when random assignment causes the type of person participating in a program to differ from the type that would participate in the program as it normally operates based on participant decisions. The description of Weikart, Bond, and McNeil (1978) suggests that because of universal participation of eligibles, this is not an issue for Perry. Substitution bias arises when members of an experimental control group gain access to close substitutes for the experimental treatment. During the pre-Head Start era of the early 1960s, there were few alternative programs to Perry, so the problem of substitution bias is unimportant for the analysis of the Perry study.

assignment protocol:

Assumption A-1. $D \sim M(R, X)$: supp $(R) \times supp(X) \rightarrow D$; $R \perp L X$, where supp(D) = D, and supp denotes support.

Let V_i represent the unobserved variables that affect outcomes for participant *i*. The vector of unobserved variables is $V = (V_i; i \in \mathcal{I})$. The assumption that unobserved variables are independent of the randomization device *R* is critical for guaranteeing that randomization produces independence between unobserved variables and treatment assignments, and can be stated as follows:

Assumption A-2. $R \perp V$.

Remark 4.1. The random variables R used to generate the randomization and the unobserved variables V are assumed to be independent. However, if initial randomization is compromised by reassignment based on X, the assignment mechanism depends on X. Thus, substantial correlation between final treatment assignments D and unobserved variables V can exist through the common dependence between X and V.

As noted in Section 2, some participants whose mothers were employed had their initial treatment status reassigned in an effort to lower program costs. One way to interpret the protocol as implemented is that the selection of reassigned participants occurred at random given working status. In this case, the assignment mechanism is based on observed variables and can be represented by M as defined in assumption A-1. In particular, conditioning on maternal working status (and other variables used to assign persons to treatment) provides a valid representation of the treatment assignment mechanism and avoids selection bias. This is the working hypothesis of our paper.

Given that many of the outcomes we study are measured some 30 years after random assignment, and a variety of post-randomization period shocks generate these outcomes, the correlation between V and the outcomes may be weak. For example, there is evidence that earnings are generated in part by a random walk with drift (see, e.g., Meghir and Pistaferri, 2004). If this is so, the correlation between the errors in the earnings equation and the errors in the assignment to treatment equation may be weak. By the proximity theorem (Fisher, 1966), the bias arising from V correlated with outcomes may be negligible.¹⁹

Each element *i* in the outcome vector *Y* takes value $Y_{i,0}$ or $Y_{i,1}$. The vectors of counterfactual outcomes are defined by $Y_d = (Y_{i,d}; i \in \mathcal{I}); d \in \{0,1\}, i \in \mathcal{I}$. Without loss of generality, assumption **A-3** postulates that outcomes $Y_{i,d}$, where $d \in \{0,1\}, i \in \mathcal{I}$, are generated by a function f:

Assumption A-3. $Y_{i,d} \equiv f(d, X_i, V_i); d \in \{0, 1\}, \forall i \in \mathcal{I}.^{20}$

Assumptions A-1, A-2, and A-3 formally characterize the Perry randomization protocol.

The Benefits of Randomization The major benefit of randomization comes from avoiding the problem of selection bias. This benefit is a direct consequence of assumptions A-1, A-2, and A-3, and can be stated as a lemma:

Lemma L-1. Under assumptions A-1, A-2, and A-3, $(Y_1, Y_0) \perp D \mid X$.

²⁰At the cost of adding new notation, we could distinguish a subset of X, Z, which does not determine M but does determine Y. In this case, we write an amended assumption:

Assumption A-3'. $Y_{i,d} = f(d, X_i, Z_i, V_i); d \in \{0, 1\}, \forall i \in \mathcal{I}.$

In addition, assumption A-2 is strengthened to the following statement:

Assumption A-2'. $R \perp (V, Z)$.

¹⁹However, if reassignment of initial treatment status was not random within the group of working mothers (say favoring those who had children with less favorable outcomes), conditioning on working status may not be sufficient to eliminate selection bias. In a companion paper, Heckman, Pinto, Shaikh, and Yavitz (2009) develop and apply a more conservative approach to bounding inference about the null hypothesis of no treatment effect where selection into treatment is based on unobserved variables correlated with outcomes, so that the assignment mechanism is described by $D \sim M(R, X, V)$. Bounding is the best that they can do because the exact rules of reassignment are unknown, and they cannot condition on V. From documentation on the Perry randomization protocol, they have a set of restrictions used to make reassignments that produce informative bounds.

In practice, conditioning on Z can be important for controlling imbalances in variables that are not used to assign treatment but that affect outcomes. For example, birth weight (a variable not used in the Perry randomization protocol) may, on average, be lower in the control group and higher in the treatment group, and birth weight may affect outcomes. In this case, a spurious treatment effect could arise in any sample due to this imbalance, and not because of the treatment itself. Such imbalance may arise from compromises in the randomization protocol. To economize on notation, we do not explicitly distinguish Z, but instead treat it as a subset of X.

Proof. Conditional on X, the argument that determines $Y_{i,d}$ for $d \in \{0,1\}$ is V, which is independent of R by assumption A-2. Thus, R is independent of (Y_0, Y_1) . Therefore, any function of R and X is also independent of (Y_0, Y_1) conditional on X. In particular, assumption A-1 states that conditional on X, treatment assignments depend only on R, so $(Y_0, Y_1) \perp D \mid X$.

Remark 4.2. Regardless of the particular type of compromise to the initial randomization protocol, Lemma L-1 is valid whenever the randomization protocol is based on observed variables X, but not on V. Assumption A-2 is a consequence of randomization. Under it, randomization provides a solution to the problem of biased selection.²¹

Remark 4.3. Lemma L-1 justifies matching as a method to correct for irregularities in the randomization protocol.

The method of matching is often criticized because the appropriate conditioning set that guarantees conditional independence is generally not known, and there is no algorithm for choosing the conditioning variables without invoking additional assumptions (e.g., exogeneity).²² For the Perry experiment, the conditioning variables X that determine the assignment to treatment are documented, even though the exact treatment assignment rule is unknown (see Weikart, Bond, and McNeil, 1978).

When samples are small and the dimensionality of covariates is large, it becomes impractical to match on all covariates. This is the "curse of dimensionality" in matching (Westat, 1981). To overcome this problem, Rosenbaum and Rubin (1983) propose propensity score matching, in which matches are made based on a propensity score, that is, the probability of being treated conditional on observed covariates. This is a one-dimensional object that

²¹Biased selection can occur in the context of a randomized experiment if treatment assignment uses information that is not available to the program evaluator and is statistically related to the potential outcomes. For example, suppose that the protocol M is based in part on an unobserved (by the economist) variable Vthat impacts Y through the $f(\cdot)$ in assumption A-3:

Assumption A-1'. M(R, X, V) : supp $(R) \times$ supp $(X) \times$ supp $(V) \rightarrow \mathcal{D}$.

²²See Heckman and Navarro (2004), Heckman and Vytlacil (2007), and Heckman (2010).

reduces the dimensionality of the matching problem at the cost of having to estimate the propensity score, which creates problems of its own.²³ Zhao (2004) shows that when sample sizes are small, as they are in the Perry data, propensity score matching performs poorly when compared with other matching estimators. Instead of matching on the propensity score, we directly condition on the matching variables using a partially linear model. A fully nonparametric approach to modeling the conditioning set is impractical in the Perry sample.

4.3 Testing the Null Hypothesis of No Treatment Effect

Our aim is to test the null hypothesis of no treatment effect. This hypothesis is equivalent to the statement that the control and treated outcome vectors share the same distribution:

Hypothesis H-1. $Y_1 \stackrel{d}{=} Y_0 \mid X$,

where $\stackrel{d}{=}$ denotes equality in distribution.

The hypothesis of no treatment effect can be restated in an equivalent form. Under Lemma L-1, hypothesis H-1 is equivalent to the following statement:

Hypothesis H-1'. $Y \perp D \mid X$.

The equivalence is demonstrated by the following argument. Let A_J denote a set in the support of a random variable J. Then

²³See Heckman, Ichimura, Smith, and Todd (1998).

 $\Pr((D,Y) \in (A_D, A_Y)|X) = E(\mathbf{1}[D \in A_D] \odot \mathbf{1}[Y \in A_Y]|X)$

(where \odot denotes a Hadamard product.²⁴) = $E(\mathbf{1}[Y \in A_Y]|D \in A_D, X) \operatorname{Pr}(D \in A_D|X)$ = $E(\mathbf{1}[(Y_1 \odot D + Y_0 \odot (1 - D)) \in A_Y]|D \in A_D, X) \operatorname{Pr}(D \in A_D|X)$ = $E(\mathbf{1}[Y_0 \in A_Y]|D \in A_D, X) \operatorname{Pr}(D \in A_D|X)$ by hypothesis H-1 = $E(\mathbf{1}[Y_0 \in A_Y]|X) \operatorname{Pr}(D \in A_D|X)$ by Lemma L-1 = $\operatorname{Pr}(Y \in A_Y|X) \operatorname{Pr}(D \in A_D|X)$,

We refer to hypotheses H-1 and H-1' interchangeably throughout this paper. If the randomization protocol is fully known, then the randomization method implies a known distribution for the treatment assignments. In this case, we can proceed in the following manner:

- 1. From knowledge of the treatment assignment rules, one can generate the distribution of D|X;
- 2. Select a statistic T(Y, D, X) with the property that larger values of the statistic provide evidence against the null hypothesis, hypothesis **H-1** (e.g., *t*-statistics, χ^2 , etc.);
- 3. Create confidence intervals for the random variable $T(Y, D, X) \mid X$ at significance level α based on the known distribution of D|X;
- 4. Reject the null hypothesis if the value of T(Y, D, X) calculated from the data does not belong to the confidence interval.

Implementing these procedures requires solving certain problems. To produce the distri-

 $^{^{24}\}mathrm{A}$ Hadamard product is an element-wise product.

bution of D|X requires precise knowledge of the ingredients of the assignment rules, which are only partially known. Alternatively, the analyst could use the asymptotic distribution of the chosen test statistic. However, given the size of the Perry sample, it seems unlikely that the distribution of T(Y, D, X) is accurately characterized by large sample distribution theory. We address these problems by using permutation-based inference that addresses the problem of small sample size in a way that allows us to simultaneously account for compromised randomization when assumptions A-1-A-3 and hypothesis H-1 are valid. Our inference is based on an *exchangeability property* that remains valid under compromised randomization.

4.4 Exchangeability and the Permutation-Based Tests

The main result of this subsection is that, under the null hypothesis, the joint distribution of outcome and treatment assignments is invariant for certain classes of permutations. We rely on this property to construct a permutation test that remains valid under compromised randomization. Permutation-based inference is often termed data-dependent because the computed p-values are conditional on the observed data. These tests are also *distributionfree* because they do not rely on assumptions about the parametric distribution from which the data are sampled. Because permutation tests give accurate p-values even when the sampling distribution is skewed, they are often used when sample sizes are small and sample statistics are unlikely to be normal. Hayes (1996) shows the advantage of permutation tests over the classical approaches for the analysis of small samples and non-normal data.

Permutation-based tests make inferences about hypothesis H-1 by exploring the invariance of the joint distribution of (Y, D) under permutations that swap the elements of the vector of treatment indicators D. We use g to index a permutation function π , where the permutation of elements of D according to π_g is represented by gD. Notationally, gD is defined as

$$gD = (\widetilde{D}_i; i \in \mathcal{I} \mid \widetilde{D}_i = D_{\pi_g(i)})$$
, where π_g is a permutation function (i.e., $\pi_g : \mathcal{I} \to \mathcal{I}$ is a bijection).

Lemma L-2. Let the permutation function $\pi_g : \mathcal{I} \to \mathcal{I}$ within each stratum of X, such that $X_i = X_{\pi_g(i)}, \ \forall i \in \mathcal{I}$. Then, under assumption **A-1**, $gD \stackrel{d}{=} D$.

Proof. $gD \sim M(R, gX)$ by construction, but gX = X by definition, so $gD \sim M(R, X)$. \Box

Remark 4.4. An important feature of the exchangeability property used in Lemma L-2 is that it relies on limited information on the randomization protocol. It is valid under compromised randomization and there is no need for a full specification of the distribution D or the assignment mechanism M.

Let \mathscr{G}_X be the set of all permutations that permute elements only within each stratum of X.²⁵ Formally,

$$\mathscr{G}_X = \{g; \ \pi_g : \mathcal{I} \to \mathcal{I} \text{ is a bijection and } X_i = X_{\pi_g(i)}, \ \forall \ i \in \mathcal{I} \}.$$

A corollary of Lemma L-2 is

$$D \stackrel{a}{=} gD \quad \forall g \in \mathscr{G}_X. \tag{1}$$

We now state and prove the following theorem.

Theorem 4.1. Let treatment assignment be characterized by assumptions A-1-A-3. Under hypothesis H-1, the joint distribution of outcomes Y and treatment assignments D is invariant under permutations $g \in \mathscr{G}_X$ of treatment assignments within strata formed by values of covariates X, that is, $(Y, D) \stackrel{d}{=} (Y, gD) \forall g \in \mathscr{G}_X$.

²⁵See Web Appendix C.3 for a formal description of restricted permutation groups.

Proof. By Lemma L-2, $D \stackrel{d}{=} gD \forall g \in \mathscr{G}_X$. But $Y \perp D \mid X$ by hypothesis H-1. Thus $(Y, D) \stackrel{d}{=} (Y, gD) \forall g \in \mathscr{G}_X$.

Theorem 4.1 is called the randomization hypothesis.²⁶ We use it to test whether $Y \perp D \mid X$. Intuitively, Theorem 4.1 states that if the randomization protocol is such that (Y, D) is invariant over the strata of X, then the absence of a treatment effect implies that the joint distribution of (Y, D) is invariant with respect to permutations of D that are restricted within strata of X.²⁷ Theorem 4.1 is a useful tool for inference about treatment effects. For example, suppose that, conditional on X (which we keep implicit), we have a test statistic T(Y, D) with the property that larger values of the statistic provide evidence against hypothesis H-1 and an associated critical value c, such that whenever T(Y, D) > c, we reject the null hypothesis. The goal of our test is to control for a Type-I error at significance level α , that is,

Pr(reject hypothesis **H-1** | hypothesis **H-1** is true)

 $= \Pr(T(Y, D) > c | \text{ hypothesis } \mathbf{H-1} \text{ is true}) \leq \alpha.$

A critical value can be computed by using the fact that as g varies in \mathscr{G}_X under the null hypothesis of no treatment effect, conditional on the sample, T(Y, gD) is uniformly distributed.²⁸ Thus, under the null, a critical value can be computed by taking the α quantile of the set $\{T(Y, gD) : g \in \mathscr{G}_X\}$. In practice, permutation tests compare a test statistic computed on the original (unpermuted) data with a distribution of test statistics computed on resamplings of that data. The measure of evidence against the randomization hypothesis, the *p*-value, is computed as the fraction of resampled data which yields a test statistic greater than that yielded by the original data. In the case of the Perry study, these resampled data sets consist of the original data with treatment and control labels permuted

²⁶See Lehmann and Romano (2005, Chapter 9).

 $^{^{27}\}mathrm{Web}$ Appendix C discusses our permutation methodology.

 $^{^{28}}$ See Lehmann and Romano (2005, Theorem 15.2.2).

across observations. As discussed below in Section 4.5, we use permutations that account for the compromised randomization, and our test statistic is the coefficient on treatment status estimated using a regression procedure due to Freedman and Lane (1983), which controls for covariate imbalances and is designed for application to permutation inference.

We use this procedure and report one-sided mid-*p*-values, which are averages between the one-sided *p*-values defined using strict and non-strict inequalities. As a concrete example of this procedure, suppose that we use a permutation test with J + 1 permutations g_j , where the first J are drawn at random from the permutation group \mathscr{G}_X and g_{J+1} is the identity permutation (corresponding to using the original sample).

Our source statistic Δ is a function of an outcome Y and permuted treatment labels $g_j D$. For each permutation, we compute a set of source statistics $\Delta^j = \Delta(Y, g_j D)$. From these, we compute the rank statistic T^j associated with each source statistic Δ^j :²⁹

$$T^{j} \equiv \frac{1}{J+1} \sum_{l=1}^{J+1} \mathbf{1} [\Delta^{j} \ge \Delta^{l}].$$
⁽²⁾

Without loss of generality, we assume that higher values of the source statistics are evidence against the null hypothesis. Working with ranks of the source statistic effectively standarizes the scale of the statistic and is an alternative to studentization (i.e., standardizing by the standard error). This procedure is called prepivoting in the literature.³⁰ The mid-p-value is

²⁹Although this step can be skipped without affecting any results for single-hypothesis testing (i.e., Δ^{j} may be used directly in calculating *p*-value), the use of rank statistics T^{j} is recommended by Romano and Wolf (2005) for the comparison of statistics in multiple-hypothesis testing.

³⁰See Beran (1988a,b). Prepivoting is defined by the transformation of a test statistic into its cumulative distribution function (cdf). The distribution is summarized by the relative ranking of the source statistics. Therefore, it is invariant to any monotonic transformation of the source statistic. Romano and Wolf (2005) note that prepivoting is useful in constructing multiple-hypothesis tests. The procedure generates a distribution of test statistics that is balanced in the sense that each prepivoting statistic has roughly the same power against alternatives. More specifically, suppose that there are no ties. After prepivoting, the marginal distribution of each rank statistic in this vector is a discrete distribution that is uniform [0, 1]. The power of the joint test of hypotheses depends only on the correlation among the prepivoting statistics, and not on their original scale (i.e., the scale of the source). The question of optimality in the choice of test statistics is only relevant to the extent that different choices change the relative ranking of the statistics. An example relevant to this paper is that the choice between tests based on difference in means across control and treatment groups or the *t*-statistic associated with the difference in means is irrelevant for permutation tests in randomized trials as both statistics produce the same rank statistics across permutations. (See Good, 2000, for a discussion.)

computed as the average of the fraction of permutation test statistics strictly greater than the unpermuted test statistic and the fraction greater than or equal to the unpermuted test statistic:

$$p \equiv \frac{1}{2(J+1)} \left(\sum_{j=1}^{J+1} \mathbf{1}[T^j \ge T^{J+1}] + \sum_{j=1}^{J+1} \mathbf{1}[T^j > T^{J+1}] \right).^{31}$$
(3)

Web Appendix C.5 shows how to use mid-*p*-values to control for Type-I error.

4.5 Accounting for Compromised Randomization

This paper solves the problem of compromised randomization under the assumption of *con*ditional exchangeability of assignments given X. A by-product of this approach is that we correct for imbalance in covariates between treatments and controls.

Conditional inference is implemented using a permutation-based test that relies on restricted classes of permutations, denoted by \mathscr{G}_X . We partition the sample into subsets, where each subset consists of participants with common background measures. Such subsets are termed *orbits* or *blocks*. Under the null hypothesis of no treatment effect, treatment and control outcomes have the same distributions within an orbit.³² Equivalently, treatment assignments D are exchangeable (therefore permutable) with respect to the outcome Y for participants who share common pre-program values X. Thus, the valid permutations $g \in \mathscr{G}_X$ swap labels *within* conditioning orbits.

We modify standard permutation methods to account for the explicit Perry randomization protocol. Features of the randomization protocol, such as identical treatment assignments for siblings, generate a distribution of treatment assignments that cannot be described (or replicated) by simple random assignment.³³

³¹Mid-*p*-values recognize the discrete nature of the test statistics.

³²The baseline variables can affect outcomes, but may (or may not) affect the distribution of assignments produced by the compromised randomization.

 $^{^{33}}$ Web Appendix C provides relevant theoretical background, as well as operational details, about implementing the permutation framework.

Conditional Inference in Small Samples Invoking conditional exchangeability decreases the number of valid permutations within X strata. The small Perry sample size prohibits very fine partitions of the available conditioning variables. In general, nonparametric conditioning in small samples introduces the serious practical problem of small or even empty permutation orbits. To circumvent this problem and obtain restricted permutation orbits of reasonable size, we assume a linear relationship between some of the baseline measures in X and the outcomes Y. We partition the data into orbits on the basis of variables that are not assumed to have a linear relationship with outcome measures. Removing the effects of some conditioning variables, we are left with larger subsets within which permutation-based inference is feasible.

More precisely, we divide the vector X into two parts: those variables $X^{[L]}$ which are assumed to have a linear relationship with Y, and variables $X^{[N]}$, whose relationship with Y is allowed to be nonparametric, $X = [X^{[L]}, X^{[N]}]$.³⁴ Linearity enters into our framework by replacing assumption A-3 with the following assumption:

Assumption A-4. $Y_{i,d} \equiv \delta_d X_i^{[L]} + f(d, X_i^{[N]}, V_i); \ d \in \{0, 1\}, \ i \in \mathcal{I}.$

Under hypothesis **H-1**, $\delta_1 = \delta_0 = \delta$ and $\tilde{Y} \equiv Y - \delta X^{[L]} = f(X^{[N]}, V)$. Using assumption **A-4**, we can rework the arguments of Section 4.4 to prove that, under the null, $\tilde{Y} \perp D \mid X^{[N]}$. Under hypothesis **A-4** and the knowledge of δ , our randomization hypothesis becomes $(\tilde{Y}, D) \stackrel{d}{=} (\tilde{Y}, gD)$ such that $g \in \mathscr{G}_{X^{[N]}}$, where $\mathscr{G}_{X^{[N]}}$ is the set of permutations that swap the participants who share the same values of covariates $X^{[N]}$. We purge the influence of $X^{[L]}$ on Y by subtracting $\delta X^{[L]}$ and can construct valid permutation tests of the null hypothesis of no treatment effect by conditioning on $X^{[N]}$. Conditioning nonparametrically on $X^{[N]}$, a smaller set of variables than X, we are able to create restricted permutation orbits that contain substantially larger numbers of observations than when we condition more finely on all of the X. In an extreme case, one could assume that all conditioning variables

³⁴Linearity is not strictly required, but we use it in our empirical work. In place of linearity, we could use a more general parametric functional form.

enter linearly, eliminate their effect on the outcome, and conduct permutations using the resulting residuals without any need to form orbits based on X.

If δ were known, we could control for the effect of $X^{[L]}$ by permuting $\tilde{Y} = Y - \delta X^{[L]}$ within the groups of participants that share the same pre-program variables $X^{[N]}$. However, δ is rarely known. We address this problem by using a regression procedure due to Freedman and Lane (1983). Under the null hypothesis, D is not an argument in the function determining Y. Our permutation approach addresses the problem raised by estimating δ by permuting the residuals from a regression of Y on $X^{[L]}$ in orbits that share the same values of $X^{[N]}$, leaving D fixed. The method regresses Y on $X^{[L]}$, then permutes the residuals from this regression according to $\mathscr{G}_{X^{[N]}}$. D is adjusted to remove the influence of $X^{[L]}$. The method then regresses the permuted residuals on adjusted D.

More precisely, define B_g as a permutation matrix associated with the permutation $g \in \mathscr{G}_{X^{[N]}}$.³⁵ The Freedman and Lane regression coefficient for permutation g is

$$\Delta_k^g \equiv (D'Q_X D)^{-1} D'Q_X B'_g Q_X Y^k; \ g \in \mathscr{G}_{X^{[N]}},\tag{4}$$

where k is the outcome index, the matrix Q_X is defined as $Q_X \equiv (\mathbf{I} - P_X)$, \mathbf{I} is the identity matrix, and

$$P_X \equiv X^{[L]}((X^{[L]})'X^{[L]})^{-1}(X^{[L]})'.$$

 P_X is a linear projection in the space generated by the columns of $X^{[L]}$, and Q_X is the projection into the orthogonal space generated by $X^{[L]}$. We use this regression coefficient as the input source statistic (Δ^j) to form the rank statistic (2) and to compute *p*-values via (3).

Expression (4) corrects for the effect of $X^{[L]}$ on both D and Y. (For notational simplicity, we henceforth suppress the k superscript.) The term $Q_X Y$ estimates \tilde{Y} . If δ were known, \tilde{Y}

³⁵A permutation matrix *B* of dimension *L* is a square matrix $B = (b_{i,j}) : i, j = 1, ..., L$, where each row and each column has a single element equal to 1 and all other elements equal to 0 within the same row or column, so $\sum_{i=1}^{L} b_{i,j} = 1$, $\sum_{j=1}^{L} b_{i,j} = 1$ for all i, j.

could be computed exactly. The term $D'Q_X$ corrects for the imbalance of $X^{[L]}$ across treatment and control groups. Without loss of generality, we can arrange the rows of (Y, D, X)so that participants that share the same values of covariates $X^{[N]}$ are adjacent. Writing the data in this fashion, B_g is a block-diagonal matrix, whose elements are themselves permutation matrices that swap elements within each stratum defined by values of $X^{[N]}$. For notational clarity, suppose that there are S of these strata indexed by $s \in S \equiv \{1, \ldots S\}$. Let the participant index set \mathcal{I} be partitioned according to these strata into S disjoint set $\{\mathcal{I}^s; s \in S\}$ so that each participant in \mathcal{I}^s has the same value of pre-program variables $X^{[N]}$. Permutations are applied within each stratum s associated with a value of $X^{[N]}$. The permutations within each stratum are conducted independently of the permutations for other strata. All within-strata permutations are generated by B_g to form Equation (4). That equation aggregates data across the strata to form Δ_k^g . The same permutation structure is applied to all outcomes in order to construct valid joint tests of multiple hypotheses. Δ_k^g plays the role of Δ^j in (2) to create our test statistic.

In a series of Monte Carlo studies, Anderson and Legendre (1999) show that the Freedman-Lane procedure generally gives the best results in terms of Type-I error and power among a number of similar permutation-based approximation methods. In another paper, Anderson and Robinson (2001) compare an exact permutation method (where δ is known) with a variety of permutation-based methods. They find that in samples of the size of Perry, the Freedman-Lane procedure generates test statistics that are distributed most like those generated by the exact method, and are in close agreement with the *p*-values from the true distribution when regression coefficients are known. Thus, for the Freedman-Lane approach, estimation error appears to create negligible problems for inference.

Interpreting Our Test Statistic To recapitulate, permutations are conducted within each stratum defined by $X^{[N]}$ for the *S* strata indexed by $s \in S \equiv \{1, \ldots, S\}$. Let D(s)be the treatment assignment vector for the subset \mathcal{I}^s defined by $D(s) \equiv (D_i; i \in \mathcal{I}^s)$. Let $\tilde{Y}(s) \equiv (\tilde{Y}_i; i \in \mathcal{I}^s)$ be the adjusted outcome vector for the subset \mathcal{I}^s . Finally, let $\mathscr{G}^s_{X^{[N]}}$ be the collection of all permutations that act on the $|\mathcal{I}^s|$ elements of the set \mathcal{I}^s of stratum s.

Note that one consequence of the conditional exchangeability property $(\tilde{Y}, D) \stackrel{d}{=} (\tilde{Y}, gD)$ for $g \in \mathscr{G}_{X^{[N]}}$ is that the distribution of a statistic within each stratum, $T(s) : (\operatorname{supp}(\tilde{Y}(s)) \times \operatorname{supp}(D(s))) \to \mathbb{R}$, is the same under permutations $g \in \mathscr{G}_{X^{[N]}}^s$ of the treatment assignment D(s). Formally, within each stratum $s \in S$,

$$T(\tilde{Y}(s), D(s)) \stackrel{d}{=} T(\tilde{Y}(s), gD(s)) \ \forall g \in \mathscr{G}^{s}_{X^{[N]}}.$$
(5)

The distribution of any statistic $T(s) = T(\tilde{Y}(s), D(s))$ (conditional on the sample) is uniform across all the values $T^g(s) = T(\tilde{Y}(s), gD(s))$, where g varies in $\mathscr{G}^s_{X^{[N]}}$.³⁶

The Freedman-Lane statistic aggregates tests across the strata. To understand how it does this, consider an approach that combines the independent statistics across strata to form an aggregate statistic,

$$T = \sum_{s=1}^{S} T(s)w(s),$$
(6)

where the weight w(s) could be, for example, $(1/\sigma(s))$ where $\sigma(s)$ is the standard error of T(s). Tests of the null hypothesis could be based on T.

To relate this statistic to the one based on Equation (4), consider the special case where there are no $X^{[L]}$ variables besides the constant term so there is no need to estimate δ . Define $D_i(s)$ as the value of D for person i in stratum $s, i = 1, \ldots, |\mathcal{I}^s|$. Likewise, $\tilde{Y}_i(s)$ is the value of \tilde{Y} for person i in stratum s. Define

$$T(s) = \frac{\sum_{i \in \mathcal{I}^s} \tilde{Y}_i(s) D_i(s)}{\sum_{i \in \mathcal{I}^s} D_i(s)} - \frac{\sum_{i \in \mathcal{I}^s} \tilde{Y}_i(s) (1 - D_i(s))}{\sum_{i \in \mathcal{I}^s} (1 - D_i(s))}.$$

We can define corresponding statistics for the permuted data.

In this special case where, in addition, the variance of $\tilde{Y}(s)$ is the same within each ³⁶See Lehmann and Romano (2005, Chapter 15) for a formal proof.

stratum ($\sigma(s) = \sigma$) and $w(s) = |\mathcal{I}^s| / \sigma |\mathcal{I}|$ (i.e., w(s) is the proportion of sample observations in stratum s), test statistic (6) generates the same inference as the Freedman-Lane regression coefficient (4) used as the source statistic for our testing procedure.

In the more general case analyzed in this paper, the Freedman-Lane procedure (4) adjusts the Y and D to remove the influence of $X^{[L]}$. Test statistic (6) would be invalid, even if we use \tilde{Y} instead of Y because it does not control for the effect of $X^{[L]}$ on $D.^{37}$ The Freedman-Lane procedure adjusts for the effect of the $X^{[L]}$, which may differ across strata.³⁸

4.6 Multiple-Hypothesis Testing: The Stepdown Algorithm

Thus far, we have considered testing a single null hypothesis. Yet there are more than 715 outcomes measured in the Perry data. We now consider the null hypothesis of no treatment effect for a set of K outcomes jointly. The complement of the joint null hypothesis is the hypothesis that there exists at least one hypothesis out of K that we reject.

Formally, let P be the distribution of the observed data, $(Y, D)|X \sim P$. We test the $|\mathcal{K}|$ set of single null hypotheses indexed by $\mathcal{K} = \{1, \ldots, K\}$ and defined by the rule

$$P \in \mathcal{P}_k \iff Y^k \perp \!\!\!\perp D | X.$$

The hypothesis we test is defined as follows:

Hypothesis H-2. $H_{\mathcal{K}}: P \in \bigcap_{k \in \mathcal{K}} \mathcal{P}_k.$

³⁷Anderson and Robinson (2001) discuss the poor performance of permutation tests that do not control for the influence of $X^{[L]}$.

³⁸The Freedman-Lane statistic is based on an OLS estimator. In the case of heteroscedasticity arising from differences in the variances of Y(s) across strata, OLS is unbiased and consistent for the treatment effect, but the conventional standard errors for OLS are biased. Asymptotic *p*-values generated using normal approximations may be misleading. Our permutation test generates valid inference by permuting data *within strata* and pooling the permuted data across strata via (4). Under the null hypothesis of no treatment effect we obtain the exact distribution of the OLS parameter conditional on the data. Thus we compute tests with the correct size. If we permuted *across* strata, we would lose this property. Whether other statistics, such as a GLS version of the Freedman-Lane statistic, would improve statistical power is still an open question. The Freedman-Lane Equation (4) is an example of a combining function in permutation statistics (Pesarin and Salmaso, 2010) applied to combine tests across strata.

The alternative hypothesis is the complement of hypothesis **H-2**. Let the unknown subset of true null hypotheses be denoted by $\mathcal{K}_P \subset \mathcal{K}$, such that $k \in \mathcal{K}_P \iff P \in \mathcal{P}_k$. Likewise we define $H_{\mathcal{K}_P} : P \in \bigcap_{k \in \mathcal{K}_P} \mathcal{P}_k$. Our goal is to test the family of null hypotheses **H-2** in a way that controls the Family-Wise Error Rate (FWER) at level α . FWER is the probability of rejecting any true null hypothesis contained in $H_{\mathcal{K}_P}$ out of the set of hypotheses $H_{\mathcal{K}}$. FWER at level α is

$$\Pr(\text{reject } H_k : k \in \mathcal{K}_P | H_{\mathcal{K}_P} \text{ is true}) \le \alpha .$$
(7)

A multiple-hypothesis testing method is said to have strong control for FWER when Equation (7) holds for any configuration of the set of true null hypotheses \mathcal{K}_P .

To generate inference using evidence from the Perry study in a robust and defensible way, we use a stepdown algorithm for multiple-hypothesis testing. The procedure begins with the null hypothesis associated with the most statistically significant statistic and then "steps down" to the null hypotheses associated with less significant statistics. The validity of this procedure follows from the analysis of Romano and Wolf (2005), who provide general results on the use of stepdown multiple-hypothesis testing procedures.

The Stepdown Algorithm Stepdown begins by considering a set of \mathcal{K} null hypotheses, where $\mathcal{K} \equiv \{1, \ldots, K\}$. Each hypothesis postulates no treatment effect of a specific outcome, that is, $H_k : Y^k \perp D | X$; $k \in \mathcal{K}$. The set \mathcal{K} of null hypotheses is associated with a block of outcomes. We adopt the mid-*p*-value p_k as the test statistic associated with each hypothesis H_k . Smaller values of the test statistic provide evidence against each null hypothesis. The first step of the stepdown procedure is a joint test of all null hypotheses in \mathcal{K} . To this end, the method uses the maximum of the set of statistics associated with hypotheses H_k , $k \in \mathcal{K}$.

The next step of the stepdown procedure compares the computed test statistic with the α -quantile of its distribution and determines whether the joint hypothesis is rejected or not. If we fail to reject the joint null hypothesis, then the algorithm stops. If we reject the null hypothesis, then we iterate and consider the joint null hypothesis that excludes the most individually statistically significant outcome — the one that is most likely to contribute to rejection of the joint null. The method steps down and is applied to a set of K - 1null hypotheses that excludes the set of hypotheses previously rejected. In each successive step, the most individually significant hypothesis — the one most likely to contribute to the significance of the joint null hypothesis — is dropped from the joint hull hypothesis, and the joint test is performed on the reduced set of hypotheses. The process iterates until only one hypothesis remains.³⁹

Summarizing, we first construct single-hypothesis p-values for each outcome in each block. We then jointly test the null hypothesis of no treatment effect for all K outcomes. After testing for this joint hypothesis, a stepdown algorithm is performed for a smaller set of K - 1 hypotheses, which excludes the most significant hypothesis among the K outcomes. The process continues for K steps. The stepdown method provides K adjusted p-values that correct each single-hypothesis p-value for the effect of multiple-hypothesis testing.

Benefits of the Stepdown Procedure Similar to traditional multiple-hypothesis testing procedures, such as the Bonferroni or Holm procedures (see, e.g., Lehmann and Romano, 2005, for a discussion of these procedures), the stepdown algorithm of Romano and Wolf (2005) exhibits strong FWER control, in contrast with the classical tests like the F or $\chi^{2,40}$ The procedure generates as many *p*-values as there are hypotheses. Thus it provides a way to determine which hypotheses are rejected. In contrast with traditional multiple-hypothesis testing procedures, the stepdown procedure is less conservative. The gain in power comes from accounting for statistical dependencies among the test statistics associated with each individual hypothesis. Lehmann and Romano (2005) and Romano and Wolf (2005) discuss the stepdown procedure in depth. Web Appendix D summarizes the literature on multiple-

 $^{^{39}}$ See Web Appendix D for details on how we implement stepdown as well as a more general and formal description of the procedure.

⁴⁰For further discussion of stepdown and its alternatives, see Westfall and Young (1993), Benjamini and Hochberg (1995), Romano and Shaikh (2004, 2006), Romano and Wolf (2005), and Benjamini, Krieger, and Yekutieli (2006).

hypothesis testing and provides a detailed description of the stepdown procedure.

4.7 The Selection of the Set of Joint Hypotheses

There is some arbitrariness in defining the blocks of hypotheses that are jointly tested in a multiple-hypothesis testing procedure. The Perry study collects information on a variety of diverse outcomes. Associated with each outcome is a single null hypothesis. A potential weakness of the multiple-hypothesis testing approach is that certain blocks of outcomes may lack interpretability. For example, one could test all hypotheses in the Perry program in a single block.⁴¹ However, it is not clear if the hypothesis "did the experiment affect any outcome, no matter how minor" is interesting. To avoid arbitrariness in selecting blocks of hypotheses, we group hypotheses into economically and substantively meaningful categories by age of participants. Income by age, education by age, health by age, test scores by age, and behavioral indices by age are treated as separate blocks. Each block is of independent interest and would be selected by economists on a priori grounds, drawing on information from previous studies on the aspect of participant behavior represented by that block. We test outcomes by age and detect pronounced life cycle effects by gender.⁴²

5 Empirical Results

We now apply our machinery to analyze the Perry data. We find large gender differences in treatment effects for different outcomes at different ages (Heckman, 2005; Schweinhart et al., 2005). We find statistically significant treatment effects for both males and females on many outcomes. These effects persist after controlling for compromised randomization and multiple-hypothesis testing.

⁴¹In addition, using large categories of closely related variables, which are statistically insignificant, increases the probability of not rejecting the null.

 $^{^{42}}$ An alternative to multiple-hypothesis testing is to assign a monetary metric to gauge the success or failure of the program. This is done in the rate of return analysis of Heckman, Moon, Pinto, Savelyev, and Yavitz (2010).

Tables 3–6 summarize the estimated effects of the Perry program on outcomes grouped by type and age of measurement.⁴³ Tables 3 and 4 report results for females, while Tables 5 and 6 are for males. The third column of each table shows the control group means for the indicated outcomes. The next three columns are the treatment effect sizes. The unconditional effect ("uncond.") is the difference in means between the treatment group and the control group. The conditional (full) effect is the coefficient on the treatment assignment variable in linear regressions. Specifically, we regress outcomes on a treatment assignment indicator and four other covariates: maternal employment, paternal presence, socio-economic status (SES) index, and Stanford-Binet IQ, all measured at the age of study entry. The conditional (partial) effect is the estimated treatment effect from a procedure using nonparametric conditioning on a variable indicating whether SES is above or below the sample median and linear conditioning for the other three covariates. This specification is used to generate the stepdown p-values reported in this paper. The next four columns are p-values, based on different procedures explained below, for testing the null hypothesis of no treatment effect for the indicated outcome. The second-to-last column, "Gender Difference-in-Difference", tests the null hypothesis of no difference in mean treatment effects between males and females. The final column gives the available observations for the indicated outcome. Small *p*-values associated with rejections of the null are bolded.

Outcomes in each block are placed in ascending order of the partially linear Freedman-Lane *p*-value, which is described below. This is the order in which the outcomes would be discarded from the joint null hypothesis in the stepdown multiple-hypothesis testing algorithm.⁴⁴ The ordering of outcomes differs in the tables for males and females. Additionally, some outcomes are reported for only one gender when insufficient observations were available for reliable testing of the hypothesis for the other gender.

⁴³Perry follow-ups were conducted at ages 19, 27, and 40. We group the outcomes by age whenever they have strong age patterns, for example, in the case of employment or income.

 $^{^{44}\}mathrm{For}$ more on the stepdown algorithm, see Section 4.6 and Web Appendix D.

Part
Females:
Outcomes.
Main
Table

Outcome	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	:50		7	7				Part Lin	7	
	20 20 2	Mean	Uncond. ^a	Cond. (Full) ^b	Cond. (Part.) ^c	Naïve ^d	Full Lin. ^e	Partial Lin. ^f	(adj.) ^g	Gender D-in-D ^h	Available Observations
Mentally Impaired?	≤ 19	0.36	-0.28	-0.29	-0.31	.008	600.	.005	.017	.337	46
Learning Disabled?	V19	0.14	-0.14	-0.15	-0.16	.009	.016	.009	.025	.029	46
Yrs. of Special Services	V1 41 (0.46	-0.26	-0.29	-0.34	.036	.013	.013	.025	.153	51
Yrs. in Disciplinary Program	<pre><19</pre>	0.36	-0.24	-0.19	-0.27	.089	.127	.074	.074	.945	46
High School Graduation	19	0.23	0.61	0.49	0.56	.000	.000	.000	.000	.003	51
Grade Point Average	19	1.53	0.89	0.88	0.95	.000	.001	.000	.001	.009	30
Highest Grade Completed	19	10.75	1.01	0.94	1.19	.007	.008	.002	.006	.052	49
# Years Held Back	≤ 19	0.41	-0.20	-0.14	-0.21	.067	.135	.097	.178	.106	46
Vocational Training Certificate	≤40	0.08	0.16	0.13	0.16	.070	.106	.107	.107	.500	51
No Health Problems		0.83	0.05	0.12	0.07	.265	.107	.137	.576	.308	49
Alive		0.92	0.04	0.04	0.06	.273	.249	.197	.675	606.	51
Vo Treat. for Illness, Past 5 Yrs.		0.59	0.05	0.14	0.10	.369	.188	.241	0690	.806	47
Vo Non-Routine Care, Past Yr.		0.00	0.04	0.02	0.03	.484	.439	.488	.896	.549	44
Vo Sick Days in Bed, Past Yr.		0.45	-0.05	-0.04	0.06	.623	.597	.529	.781	.412	47
No Doctors for Illness, Past Yr.		0.54	-0.02	-0.01	-0.05	.559	.539	.549	.549	.609	49
Vo Tobacco Use	27	0.41	0.11	0.08	0.08	.208	.348	.298	.598	.965	47
Infrequent Alcohol Use	27	0.67	0.17	0.07	0.12	.103	.336	.374	.587	.924	45
Routine Annual Health Exam	27	0.86	-0.06	-0.09	-0.05	.684	.751	.727	.727	.867	47
Ias Anv Children	< 19	0.52	-0.12	-0.05	-0.07	.218	.419	.328	.601		48
# Out-of-Wedlock Births	≤ 40	2.52	-0.29	0.51	0.05	.652	.257	.402	.402		42
⊭ Non-Juv. Arrests	<27	1.88	-1.60	-2.22	-2.14	.016	.003	.003	.005	.571	51
Any Non-Juv. Arrests	≤ 27	0.35	-0.15	-0.18	-0.14	.148	.122	.125	.125	.440	51
# Total Arrests	≤40	4.85	-2.65	-2.88	-2.77	.028	.037	.041	.088	.566	51
# Total Charges	≤ 40	4.92	2.68	2.81	2.81	.030	.037	.042	.088	.637	51
⊭ Non-Juv. Arrests	≤ 40	4.42	-2.26	-2.62	-2.45	.044	.046	.051	.102	.458	51
# Misd. Arrests	<40	4.00	-1.88	-2.19	-2.02	.078	.078	.085	.160	.549	51
Total Crime Cost ⁱ	≤ 40	293.50	-271.33	-381.03	-381.03	.013	.108	060.	060.	.858	51
Any Arrests	≤ 40	0.65	-0.09	-0.11	-0.13	.181	.280	.239	.310	.824	51
Any Charges	140	0.65	0.09	0.13	0.13	.181	.280	.239	.310	.799	51
Any Non-Juv. Arrests Nuv Misch Arrests	<pre></pre>	0.54 0.54	-0.02	-0.02	-0.02	.351 351	.541	.520	.520	.463	51
	Alive contracture for Illness, Past 5 Yrs. No Treat, for Illness, Past 5 Yrs. No Sick Days in Bed, Past Yr. No Doctors for Illness, Past Yr. No Tobacco Use Infrequent Alcohol Use Infrequent Alcohol Use Routine Annual Health Exam Has Any Children # Out-of-Wedlock Births # Non-Juv. Arrests Any Non-Juv. Arrests # Total Arrests # Total Arrests # Misd. Arrests Any Arrests Any Arrests Any Arrests Any Non-Juv. Arrests	Allve40No Treat, for Illness, Past 5 Yrs.27No Non-Koutine Care, Past Yr.27No Sick Days in Bed, Past Yr.27No Sick Days in Bed, Past Yr.27No Doctors for Illness, Past Yr.19No Tobacco Use27Infrequent Alcohol Use27Infrequent Alcohol Use27Routine Annual Health Exam27Has Any Children27Has Any Children27Any Non-Juv. Arrests ≤ 40 # Total Arrests ≤ 240 # Misd. Arrests ≤ 40 # Misd. Arrests ≤ 40 Any Arrests ≤ 40 Any Arrests ≤ 40 Any Non-Juv. Arrests ≤ 40 Any Arrests ≤ 40 Any Non-Juv. Arrests ≤ 40 Any Arrests ≤ 40 Any Non-Juv. Arrests ≤ 40		$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$

(c) Conditional treatment effect as in the previous column except that SES is replaced with an indicator for SES above/below the median, so that the corresponding *p*-value Notes: Monetary values adjusted to thousands of year-2006 dollars using annual national CPI. *p*-values below 0.1 are in bold. (a) Unconditional difference in means between the treatment and control groups; (b) Conditional treatment effect with linear covariates Stanford-Binet IQ, socio-economic status index (SES), maternal employment, father's is computed in the column "Partial Lin." This specification generates *p*-values used in the stepdown procedure; (d) One-sided *p*-values for the hypothesis of no treatment effect based on conditional permutation inference, without orbit restrictions or linear covariates — estimated effect size in the "Uncond." column; (e) One-sided *p*-values employment, paternal presence, socio-economic status index (SES), and Stanford-Binet IQ) — estimated effect size in the "Conditional Effect" column; (f) One-sided *p*-values for the hypothesis of no treatment effect based on the Freedman-Lane procedure, using the linear covariates maternal employment, paternal presence, and Stanford-Binet IQ, p-values from the previous column, adjusted for multiple inference using the stepdown procedure; (h) Two-sided p-value for the null hypothesis of no gender difference in mean treatment effects, tested using mean differences between treatments and controls using the conditioning and orbit restriction setup described in footnote (e); (i) Total crime of the Midwest, police and court costs are based on historical Michigan unit costs, and the victimization cost of fatal crime takes into account the statistical value of life (see for the hypothesis of no treatment effect based on the Freedman-Lane procedure, without restricting permutation orbits and assuming linearity in all covariates (maternal and restricting permutation orbits within strata formed by Socio-economic Status index (SES) being above or below the sample median and permuting siblings as a block; (g) costs include victimization, police, justice, and incarceration costs, where victimizations are estimated from arrest records for each type of crime using data from urban areas presence at study entry - this is also the effect for the Freedman-Lane procedure under a full linearity assumption, whose respective *p*-value is computed in column "Full Lin.", Heckman, Moon, Pinto, Savelyev, and Yavitz, 2010, for details)

	Outcome	Age	Ctl.	Effect	st Cond.	Cond.	۳ : :	<i>p</i> -vε Full	p -values ll Partial	Part. Lin.	Gender	Available
		_	Mean	Uncond."	(Full) ^b	(Part.) ^c	Naive	$Lin.^{e}$	$\operatorname{Lin.}^{\mathrm{f}}$	(adj.) ^g	$D-in-D^h$	Observations
tuə	No Job in Past Year Jobless Months in Past 2 Yrs. Current Employment	19 19 19	$\begin{array}{c} 0.58 \\ 10.42 \\ 0.15 \end{array}$	-0.34 -5.20 0.29	-0.37 -5.47 0.23	-0.38 -6.82 0.27	.006 .054 .023	.007 .099 .045	.003 .020 .032	.007 .036 .032	.009 .102 .373	51 42 51
utolqn	No Job in Past Year Current Employment Jobless Months in Past 2 Yrs.	27 27 27	$\begin{array}{c} 0.54 \\ 0.55 \\ 10.45 \end{array}$	-0.29 0.25 -4.21	-0.25 0.18 -2.14	-0.30 0.28 -4.23	.017 .036 .077	.058 .096 .285	.037 .042 .165	.071 .063 .165	.157 .220 .908	48 47 47
тЭ	No Job in Past Year Jobless Months in Past 2 Yrs. Current Employment	40 40	0.41 5.05 0.82	-0.25 -1.05 0.02	-0.22 1.05 -0.08	-0.24 -0.60 -0.01	.032 .343 .419	.092 .654 .727	.056 .528 .615	.111 .627 .615	.464 .573 .395	47 46 46
, is	Monthly Earn., Current Job	19	2.08	-0.61	-0.47	-0.51	.750	.701	.725		.677	15
Zuinı	Monthly Earn., Current Job Yearly Earn., Current Job	27 27	$1.13 \\ 15.45$	0.69 4.60	$0.48 \\ 2.18$	0.64 4.00	.050	.144 .339	.109 .277	.139 .277	.752 .873	47
вa	Yearly Earn., Current Job Monthly Earn., Current Job	40	19.85 1.85	4.35 0.21	$4.46 \\ 0.27$	5.27 0.38	.328	.272 .316	.224 .261	.274 .261	.755	46 46
i.qn	No Job in Past Year Jobless Months in Past 2 Yrs. Current Employment Monthly Earn., Current Job	19 19 19	$\begin{array}{c} 0.58 \\ 10.42 \\ 0.15 \\ 2.08 \end{array}$	-0.34 -5.20 0.29 -0.61	-0.37 -5.47 0.23 -0.47	-0.38 -6.82 0.27 -0.51	.006 .054 .023	.007 .099 .045	.003 .020 .725	.010 .056 .064 .725	.009 .102 .373 .677	51 42 15 15
13 % syniny	No Job in Past Year Current Employment Monthly Barn, Current Job Jobless Months in Past 2 Yrs. Yearly Earn, Current Job	27 27 27 27 27	$\begin{array}{c} 0.54 \\ 0.55 \\ 1.13 \\ 10.45 \\ 15.45 \end{array}$	-0.29 0.25 0.69 4.60	-0.25 0.18 0.48 -2.14 2.18	-0.30 0.28 0.64 4.23	.017 .036 .036 .050 .077	.058 .096 .144 .285 .339	.037 .042 .109 .165 .277	.094 .094 .188 .241 .277	.157 .220 .968 .873	44 44 44 44 44 44 44 44 44 44 44 44 44
មេភ្ញ	No Job in Past Year Yearly Earn., Current Job Monthly Earn., Current Job Jobless Months in Past 2 Yrs. Current Employment	40 40 40 40 40	$\begin{array}{c c} 0.41 \\ 19.85 \\ 1.85 \\ 5.05 \\ 0.82 \end{array}$	-0.25 4.35 0.21 -1.05 0.02	-0.22 4.46 0.27 1.05 -0.08	-0.24 5.27 0.38 -0.60 -0.01	.032 .251 .328 .343 .419	.092 .272 .316 .654 .727	.056 .224 .261 .528 .615	.156 .423 .440 .627 .615	.464 .755 .708 .573 .395	47 46 46 46
	Savings Account Car Ownership Checking Account	27 27 27	0.45 0.59 0.27	$\begin{array}{c} 0.27 \\ 0.13 \\ 0.01 \end{array}$	$\begin{array}{c} 0.23 \\ 0.12 \\ -0.03 \end{array}$	0.26 0.18 0.00	.036 .164 .472	.0 87 .221 .586	.051 .147 .472	.132 .250 .472	.128 .887 .777	47 47
əimonoə	Credit Card Checking Account Car Ownership Savings Account	40 40 40 40	0.50 0.50 0.77 0.73	0.04 0.08 0.06 0.06	0.06 0.04 0.03 -0.08	0.11 0.12 0.11 0.05	.425 .321 .280 .309	.355 .413 .409 .722	.233 .237 .257 .516	.483 .450 .394 .516	.737 .675 .157 .071	46 46 46 46
Э	Ever on Welfare > 30 Mos. on Welfare # Months on Welfare Never on Welfare Never on Welfare (Self Rep.)	$\begin{array}{c c}18-27\\18-27\\18-27\\16-40\\26-40\end{array}$	$\begin{array}{c} 0.82 \\ 0.55 \\ 51.23 \\ 0.92 \\ 0.41 \end{array}$	-0.34 -0.27 -21.51 -0.16 0.09	-0.21 -0.18 -11.39 -0.13 0.14	-0.27 -0.25 -21.58 -0.12 0.06	.009 .036 .060 .110 .759	.084 .152 .241 .129 .787	.049 .072 .120 .132 .664	.154 .187 .265 .221 .664	.074 .087 .122 .970 .118	47 47 51 46

control groups; (b) Conditional treatment effect with linear covariates Stanford-Binet IQ, socio-economic status index (SES), maternal employment, father's presence at study entry — this is also the effect for the Freedman-Lane procedure under a full linearity assumption, whose respective *p*-value is computed in column "Full Lin."; (c) Conditional treatment effect as in the previous column except that SES is replaced with an indicator for SES above/below the median, so that the corresponding *p*-value is computed in the column "Full Lin." This specification generates *p*-values used in the stepdown procedure; (d) One-sided *p*-values for the hypothesis of no treatment effect based on conditional permutation inference, without orbit restrictions or linear covariates — estimated effect size in the "Uncond." column; (e) One-sided *p*-values for the hypothesis of no treatment effect based on the Freedman-Lane procedure, without restricting permutation orbits and assuming linearity in all covariates (maternal employment, paternal presence, socio-economic status index (SES), and Stanford-Binet IQ) — estimated effect size in the "Conditional Effect" column; (f) One-sided *p*-values for the hypothesis of no treatment effect based on the Freedman-Lane procedure, using the linear covariates maternal employment, paternal presence, and Stanford-Binet IQ, and restricting permutation orbits within strata formed by Socio-economic Status index (SES) being above or below the sample median and permuting siblings as a block; (g) *p*-values from the previous column, adjusted for multiple inference using the stepdown procedure; (h) Two-sided *p*-value for the null hypothesis of no gender difference in mean treatment effects, tested using mean differences between treatments and controls using the conditioning and orbit restriction setup described in footnote (e); Monetary values adjusted to thousands of year-2006 dollars using annual national CPI. *p*-values below 0.1 are in bold. (a) Unconditional difference in means between the tria standard species (b) Conditional treatment effect with linear covariates Stanford-Bine IQ, socio-economic status index (SES), maternal employament, father's presence at study entry — this is also the effect for the Freadman-Lane procedure under a full linearity assumption, whose respective *p*-value is computed in column "Full Lin."; (c) Conditional treatment effect with an indicator for SES above/below the median, so that the corresponding *p*-value is computed in the column (c) Conditional treatment effect as in the previous column except that SES is replaced with an indicator for SES above/below the median, so that the corresponding *p*-value is computed in the column (c). and Stanford-Binet IQ, and restricting permutation orbits within strata formed by Socio-economic Status index (SES) being above or below the sample median and permuting siblings as a block; (g) *p*-values from the previous column, adjusted for multiple inference using the stepdown procedure; (h) Two-sided *p*-value for the null hypothesis of no gender difference in mean treatment effects, tested using the stepdown procedure; (h) Two-sided *p*-value for the null hypothesis of no gender difference in mean treatment effects, tested using the conditioning and orbit restriction setup described in footnote (e); (i) Age-19 measures are conditional on at least some earnings during the period specified — observations with zero earnings are omitted in computing means and regressions. (a) Unconditional difference in means between the treatment and orbit restrictions or linear covariates — estimated effect size in the "Uncond." column; (e) One-sided *p*-values for the hypothesis of no treatment effect based on the Freedman-Lane procedure, without restricting permutation orbits and assuming linearity in all covariates (maternal employment, paternal presence, socio-economic status index (SES), and Stanford-Binet IQ) — estimated effect size in the ¹Partial Lin." This specification generates *p*-values used in the stepdown procedure; (d) One-sided *p*-values for the hypothesis of no treatment effect based on conditional permutation inference, without "Conditional Effect" column; (f) One-sided p-values for the hypothesis of no treatment effect based on the Freedman-Lane procedure, using the linear covariates maternal employment, paternal presence, Monetary values adjusted to thousands of year-2006 dollars using annual national CPI. p-values below 0.1 are in bold. Notes:

Table 5: Main Outcomes, Males: Part 1

<pre></pre>	0.42 -0.12 -0.26 0.46 -0.14 -0.10
	-0.04 0.08
00	33 0.06 0.06 51 -0.03 0.00
0	0.08
.0	0.05
000	0.10
0.50	0.07
0.17 0.95	-0.03 -0.02
0.58	
0.46	
0.74	
5.36	
2.33	
0.72	
0.92	
10.05	
0.87	
8.46	0.18 0.12 -0.04 -1.12 -1.12 -0.00 0.00 -0.14 -0.14 -0.13
11.72	
12.41 3.26	
3.08	
775.90	
0.62	0.18 0.12 -0.04 -2.33 -1.12 -0.02 0.00 0.00 0.00 -0.14 -0.14 -0.15 -0.15 -0.13 -0.13 -0.13 -0.13 -0.13 -1.12 -1.12 -1.12 -1.12 -1.12 -1.12 -0.03 -0.03 -0.00 -0.03 -0.03 -0.15 -0.03 -0.15 -0.05 -0.15 -0.05 -0.00 -0.15 -0.05 -0.15 -0.05 -0.15 -1.12 -0.15 -1.12 -0.15 -0.115 -0.115 -1.12 -1.12 -0.115 -1.12 -0.115 -1.12 -1.12 -0.115 -1.12 -1
0.23 0.95	0.18 0.12 0.12 -0.04 -1.12 -1.12 -0.13 -0.14 -0.16 -0.13 -0.13 -0.13 -0.13 -0.13 -0.14 -0.13 -0.14 -0.13 -0.15 -1.12 -0.13 -0.13 -0.13 -0.13 -0.13 -0.13 -0.13 -0.13 -0.13 -0.13 -0.13 -0.13 -0.13 -0.13 -0.13 -0.13 -0.13 -0.14 -0.13 -0.14 -0.13 -0.14 -0.13 -0.14 -0.13 -1.14 -0.13 -1.14 -0.13 -1.14 -1.14 -0.13 -1.14

(c) Conditional treatment effect as in the previous column except that SES is replaced with an indicator for SES above/below the median, so that the corresponding *p*-value is computed in the column "Partial Lin." This specification generates *p*-values used in the stepdown procedure; (d) One-sided *p*-values for the hypothesis of no treatment effect based on conditional permutation inference, without orbit restrictions or linear covariates — estimated effect size in the "Uncond." column; (e) One-sided *p*-values Notes: Monetary values adjusted to thousands of year-2006 dollars using annual national CPI. *p*-values below 0.1 are in bold. (a) Unconditional difference in means between the treatment and control groups; (b) Conditional treatment effect with linear covariates Stanford-Binet IQ, socio-economic status index (SES), maternal employment, father's for the hypothesis of no treatment effect based on the Freedman-Lane procedure, without restricting permutation orbits and assuming linearity in all covariates (maternal — estimated effect size in the "Conditional Effect" column; (f) One-sided p-values *p*-values from the previous column, adjusted for multiple inference using the stepdown procedure; (h) Two-sided *p*-value for the null hypothesis of no gender difference in mean treatment effects, tested using mean differences between treatments and controls using the conditioning and orbit restriction setup described in footnote (e); (i) Non-victimless crimes are those associated with victimization costs: murder, rape, robbery, assault, burglary, larceny, and motor vehicle theft (see Heckman, Moon, Pinto, Savelyev, and Yavitz, 2010, for details); (j) Total crime costs include victimization, police, justice, and incarceration costs, where victimizations are estimated from arrest records for each type of crime using data from urban areas of the Midwest, police and court costs are based on historical Michigan unit costs, and the victimization cost of fatal crime takes into account presence at study entry — this is also the effect for the Freedman-Lane procedure under a full linearity assumption, whose respective *p*-value is computed in column "Full Lin."; for the hypothesis of no treatment effect based on the Freedman-Lane procedure, using the linear covariates maternal employment, paternal presence, and Stanford-Binet IQ, and restricting permutation orbits within strata formed by Socio-economic Status index (SES) being above or below the sample median and permuting siblings as a block; (g) the statistical value of life (see Heckman, Moon, Pinto, Savelyev, and Yavitz, 2010, for details). employment, paternal presence, socio-economic status index (SES), and Stanford-Binet IQ)

Available Observations	72 70 72	69 72 69	66 66 72	30	68 66	66 66	72 30 72	68 69 66 69 69 69	66 72 66 66	02 02	66 66 66	72 66 66 66
Gender D-in-D ^h	.373 .102 .009	.908 .157 .220	.395 .573 .464	.677	.752 .873	.755 .708	.373 .677 .102 .009	. 752 . 908 . 873 . 157 . 220 . 395	.573 .464 .755 .708	.887 .128 .777	.071 .157 .737 .675	.970 .118 .0 87 .122 .0 74
Part. Lin. (adj.) ^g	.196 .841 .857	.065 .294 .219	.024 .026 .068		.018 .186	.203 .195	.279 .736 .841 .857	.037 .084 .360 .294 .219 .035	.045 .137 .203 .195	.152 .610 .575	.003 .004 .327 .491	.104 .147 .619 .646 .590
<i>p</i> -values Partial Lin. ^f	.103 .781 .857	.033 .192 .219	.011 .018 .068	.442	.011 .186	.150 .195	.103 .442 .781 .857	.011 .033 .186 .192 .219 .011	.018 .068 .150 .195	.059 .397 .575	.001 .002 .206 .491	.028 .051 .517 .590
Full Lin. ^e	.144 .763 .827	.057 .295 .251	.011 .040 .123	.408	.014 .248	.270 .277	.144 .408 .763 .827	.014 .057 .248 .295 .251 .011	.040 .123 .270 .277	.072 .425 .610	.002 .003 .558	.086 .075 .482 .566 .635
Naïve ^d	.101 .784 .924	.059 .260 .367	.0 59 .082 .249	.591	.017 .227	.147 .224	.101 .591 .784 .924	.017 .059 .227 .260 .367 .059	.082 .249 .147 .224	.089 .555 .591	.002 .004 .180 .463	.101 .058 .571 .563 .697
Cond. (Part.) ^c	$\begin{array}{c} 0.16 \\ 1.50 \\ 0.10 \end{array}$	-4.50 -0.09 0.10	0.29 -5.17 -0.17	0.13	1.01 4.38	7.02 0.55	0.16 0.13 1.50 0.10	1.01 -4.50 4.38 -0.09 0.10	-5.17 -0.17 7.02 0.55	0.19 0.04 -0.02	0.38 0.35 0.10 0.01	-0.19 -0.20 -0.01 0.37 0.03
Effect Cond. (Full) ^b	$\begin{array}{c} 0.13 \\ 1.31 \\ 0.09 \end{array}$	-4.09 -0.07 0.09	0.29 -4.59 -0.15	0.09	0.99 3.67	$4.62 \\ 0.44$	$\begin{array}{c} 0.13 \\ 0.09 \\ 1.31 \\ 0.09 \end{array}$	$\begin{array}{c} 0.99\\ -4.09\\ 3.67\\ -0.07\\ 0.09\\ 0.29 \end{array}$	-4.59 -0.15 4.62 0.44	$\begin{array}{c} 0.18 \\ 0.03 \\ -0.02 \end{array}$	$\begin{array}{c} 0.36 \\ 0.32 \\ 0.08 \\ -0.01 \end{array}$	-0.17 -0.18 -0.02 -0.14 0.02
Uncond. ^a	$\begin{array}{c} 0.14\\ 1.47\\ 0.11\end{array}$	-3.66 -0.07 0.04	0.20 -3.52 -0.10	-0.16	0.88 3.50	$7.17 \\ 0.50$	$\begin{array}{c} 0.14 \\ -0.16 \\ 1.47 \\ 0.11 \end{array}$	0.88 -3.66 3.50 -0.07 0.04 0.04	-3.52 -0.10 7.17 0.50	0.15 -0.01 -0.04	$\begin{array}{c} 0.37\\ 0.30\\ 0.11\\ 0.01\end{array}$	-0.15 -0.18 -0.01 0.59 0.06
Ctl. Mean	0.41 3.82 0.13	$8.79 \\ 0.31 \\ 0.56$	$\begin{array}{c} 0.50 \\ 10.75 \\ 0.46 \end{array}$	2.74	$1.43 \\ 21.51$	$24.23 \\ 2.11$	$\begin{array}{c} 0.41 \\ 2.74 \\ 3.82 \\ 0.13 \end{array}$	$\begin{array}{c} 1.43\\ 8.79\\ 8.79\\ 21.51\\ 0.31\\ 0.56\\ 0.50\end{array}$	$10.75 \\ 0.46 \\ 24.23 \\ 2.11$	$\begin{array}{c} 0.59 \\ 0.46 \\ 0.23 \end{array}$	0.36 0.50 0.36 0.39	$\begin{array}{c} 0.82\\ 0.38\\ 0.08\\ 6.84\\ 0.26\end{array}$
Age	19 19 19	27 27 27	40 40	19	27 27	40 40	19 19 19	27 27 27 27 40	40 40 40	27 27 27	40 40 40	$\begin{array}{c} 16-40\\ 26-40\\ 18-27\\ 18-27\\ 18-27\\ 18-27\end{array}$
Outcome	Current Employment Jobless Months in Past 2 Yrs. No Job in Past Year	Jobless Months in Past 2 Yrs. No Job in Past Year Current Employment	Current Employment Jobless Months in Past 2 Yrs. No Job in Past Year	Monthly Earn., Current Job	Monthly Earn., Current Job Yearly Earn., Current Job	Yearly Earn., Current Job Monthly Earn., Current Job	Current Employment Monthly Earn., Current Job Jobless Months in Past 2 Yrs. No Job in Past Year	Monthly Earn, Current Job Jobless Months in Past 2 Yrs. Yearly Earn, Current Job No Job in Past Year Current Employment Current Employment	Jobless Months in Past 2 Yrs. No Job in Past Year Yearly Earn., Current Job Monthly Earn., Current Job	Car Ownership Savings Account Checking Account	Savings Account Car Ownership Credit Card Checking Account	Never on Welfare Never on Welfare (Self Rep.) > 30 Mos. on Welfare # Months on Welfare Ever on Welfare
	зuə	myolqm	чЭ	is	guin1.	EЭ	ⁱ .qm	1.J 28 syninreA	ī I	I	oimonoo2	I I

the treatment and control groups; (b) Conditional treatment effect with linear covariates Stanford-Binet IQ, socio-economic status index (SES), maternal employment, father's (c) Conditional treatment effect as in the previous column except that SES is replaced with an indicator for SES above/below the median, so that the corresponding *p*-value effect based on conditional permutation inference, without orbit restrictions or linear covariates — estimated effect size in the "Uncond" column; (e) One-sided p-values employment, paternal presence, socio-economic status index (SES), and Stanford-Binet IQ) — estimated effect size in the "Conditional Effect" column; (f) One-sided *p*-values and restricting permutation orbits within strata formed by Socio-economic Status index (SES) being above or below the sample median and permuting siblings as a block; (g) p-values from the previous column, adjusted for multiple inference using the stepdown procedure; (h) Two-sided p-value for the null hypothesis of no gender difference in mean treatment effects, tested using mean differences between treatments and controls using the conditioning and orbit restriction setup described in footnote (e); (i) Age-19 Notes: Monetary values adjusted to thousands of year-2006 dollars using annual national CPI. *p*-values below 0.1 are in bold. (a) Unconditional difference in means between is computed in the column "Partial Lin." This specification generates *p*-values used in the stepdown procedure; (d) One-sided *p*-values for the hypothesis of no treatment for the hypothesis of no treatment effect based on the Freedman-Lane procedure, without restricting permutation orbits and assuming linearity in all covariates (maternal for the hypothesis of no treatment effect based on the Freedman-Lane procedure, using the linear covariates maternal employment, paternal presence, and Stanford-Binet IQ, presence at study entry — this is also the effect for the Freedman-Lane procedure under a full linearity assumption, whose respective p-value is computed in column "Full Lin."; measures are conditional on at least some earnings during the period specified — observations with zero earnings are omitted in computing means and regressions. **Single** *p***-values** Tables 3–6 show four varieties of *p*-values for testing the null hypothesis of no treatment effect. The first such value, labeled "Naïve", is based on a simple permutation test of the hypothesis of no difference in means between treatment and control groups. This test uses no conditioning, imposes no restrictions on the permutation group, and does not account for imbalances or the compromised Perry randomization. These naïve *p*-values are very close to their asymptotic versions. For evidence on this point, see Web Appendix E.

The next three *p*-values are based on variants of a procedure due to Freedman and Lane (1983) for combining regression with permutation testing for admissible permutation groups. The first Freedman-Lane p-value, labeled "Full Linearity", tests the significance of the treatment effect, adjusting outcomes using linear regression with four covariates: maternal employment, paternal presence, SES, and Stanford-Binet IQ, all measured at study entry.⁴⁵ The second Freedman-Lane *p*-value, labeled "Partial Linearity", allows for a nonparametric relationship between the SES index and outcomes while continuing to assume a linear relationship for the other three covariates. This nonparametric conditioning on SES is achieved by restricting the orbits of the permutations used in the test. Exchangeability of treatment assignments between observations is assumed only on subsamples with similar values of the SES index (specifically, whether subjects fall above or below the sample median). In addition, the permutation distribution for the partially linear *p*-values permute siblings as a block. Admissible permutations do not assign different siblings to different treatment and control statuses. These two modifications account for the compromised randomization of the Perry study.⁴⁶ The third *p*-value for the Freedman-Lane procedure incorporates an adjustment for multiple-hypothesis testing using the stepdown algorithm described below.

Stepdown *p***-values and Multiple-Hypothesis Testing** We divide outcomes into blocks for multiple-hypothesis testing by type of outcome, similarities on the type of measure, and

⁴⁵Note that these are the same four used to produce the conditional effect size previously described.

⁴⁶Partial linearity is a valid assumption if full linearity is a valid assumption, although the converse need not necessarily hold since a nonparametric approach is less restrictive than a linear parametric approach.

age if there is an obvious age pattern.⁴⁷ In Tables 3–6, these blocks are delineated by horizontal lines. Within each block, the "Partially Linear (Adjusted)" p-value is the set of p-values obtained from the partially linear model adjusted for multiple-hypothesis testing using the stepdown algorithm. The adjusted p-value in each row corresponds to a joint hypothesis test of the indicated outcome and the outcomes within each block.

The first row of each block constitutes a joint test of the null hypothesis of no treatment effect for any of the outcomes in that block. Each successive row eliminates one outcome from the joint null hypothesis. This stepwise ordering is the reason why we report outcomes placed in ascending order of their *p*-values. The stepdown-adjusted *p*-values are based on these values, and the most individually significant remaining outcome is removed from the joint null hypothesis at each successive step.

Statistics We use the mid-*p*-value statistics based on the Freedman-Lane coefficient Δ_k^g for treatment status *D*. All *p*-values are computed using 30,000 draws under the relevant permutation procedure. All inference is based on one-sided *p*-values under the assumption that treatment is not harmful. An exception is the test for differences in treatment effects by gender, which are based on two-sided *p*-values.

Main Results Tables 3–6 show many statistically significant treatment effects and gender differences that survive multiple-hypothesis testing. In summary, females show strong effects for educational outcomes, early employment, and other early economic outcomes, as well as reduced numbers of arrests. Males show strong effects on a number of outcomes, demonstrating a substantially reduced number of arrests and lower probability of imprisonment, as well as strong effects on earnings at age 27, employment at age 40, and other economic outcomes recorded at age 40.

A principal contribution of this paper is to *simultaneously* tackle the statistical challenges

⁴⁷Education, health, family composition, criminal behavior, employment status, earnings, and general economic activities are the categories of variables on which blocks are selected on a priori grounds.

posed by the problems of small sample size, imbalance in the covariates, and compromised randomization. In doing so, we find substantial differences in inference between the testing procedures that use naïve p-values versus the Freedman-Lane p-values which correct for the compromised nature of the randomization protocol. The rejection rate when correcting for these problems is often higher compared with what is obtained from procedures that do not make such corrections, sharpening the evidence for treatment effects from the Perry program. This pattern is largely found in the p-values for males. This is evidenced by increasing statistical significance of treatment effects moving from "Naïve" to "Full Linearity" and from "Full Linearity" to "Partial Linearity". In several cases, outcomes that are statistically insignificant at a 10% level using naïve p-values are shown to be statistically significant using p-values for "Current Employment" at age 40 for males or "Non-juvenile Arrests" at age 27 for females.

Schooling Within the group of hypotheses for education, the only statistically significant treatment effect for males is the effect associated with being classified as mentally impaired through age 19 (Table 5). We fail to reject the overall joint null hypotheses for both school achievement and for lifetime educational outcomes. However, as Table 3 shows, there are strong treatment effects for females on high school GPA, graduation, highest grade completed, mental impairment, learning disabilities, and so on. The hypothesis of no difference between sexes in schooling outcomes is rejected for the outcomes of highest grade completed, GPA, high school graduation, and the presence of a learning disability. The unimpressive education results for males, however, do not necessarily mean that the pattern would be reproduced if the program were replicated today. We discuss this point in Section 6.⁴⁸ We discuss the effects of the intervention on cognitive test scores in Web Appendix G. Heckman, Malofeeva, Pinto, and Savelyev (2010) discuss the impact of the Perry program on noncognitive skills. They decompose treatments effects into effects due to cognitive and noncognitive

⁴⁸We present a more extensive discussion of this point in Web Appendix I.

enhancements of the program.

Employment and Earnings Results for employment and earnings are displayed in Table 4 for females and Table 6 for males. The treatment effects in these outcomes exhibit gender differences and a distinctive age pattern. For females, we observe statistically significant employment effects in the overall joint null hypotheses at ages 19 and 27. Only one outcome does not survive stepdown adjustment: "Jobless Months in Past 2 Years" at age 27. At age 40, however, there are no statistically significant earnings effects for females considered as individual outcomes or in sets of joint null hypotheses by age. For males, we observe no significant employment effects at age 19. We reject the overall joint null hypotheses of no difference in employment outcomes at ages 27 and 40. We also reject the null hypotheses of no treatment effect on age-40 employment outcomes individually. When male earnings outcomes alone are considered, we reject only the overall joint null hypothesis at age 27. However, when earnings are considered together with employment, we reject both the overall age-27 and age-40 joint null hypotheses.

Economic Activity Tests for other economic outcomes, shown in Tables 4 and 6, reinforce the conclusions drawn from the analysis of employment outcomes above. Both treated males and females are generally more likely to have savings accounts and own cars at the same ages that they are more likely to be employed. The effects on welfare dependence are strong for males when considered through age 40, but weak when considered only through age 27; the converse is true for females.

Criminal Activity Tables 3 and 5 show strong treatment effects on criminal activities for both genders. Males are arrested far more frequently than females, and, on average, male crimes tend to be more serious. There are no statistically significant gender differences in treatment effects for comparable crime outcomes. By age 27, control females were arrested 1.88 times on average during adulthood, including 0.27 felony arrests, while the correspond-

ing figures for control males are 5.36 and 2.33.⁴⁹ In addition, treated males are significantly less likely to be in prison at age 40 than their control counterparts.⁵⁰ Figure 4 shows cumulative distribution functions for charges cited at all arrests through age 40 for the male subsample. Figure 4a includes all types of charges, while Figure 4b includes only charges with nonzero victim costs. The latter category of charges is relevant because the costs of criminal victimization resulting from crimes committed by the Perry subjects play a key role in determining the economic return to the Perry Preschool Program. This is reflected in the statistical significance of estimated differences in total crime costs between treated and untreated groups at the 10% level based on the Freedman-Lane procedure using the partially linear model for both males and females. Total crime costs include victimization, police, justice, and incarceration costs. Victimizations are estimated from arrest records for each type of crime using data from urban areas of the Midwest. Police and court costs are based on historical Michigan unit costs, and the victimization cost of fatal crime takes into account the statistical value of life.⁵¹ We reject the overall joint null hypotheses for the number of arrests for both males and females at age 27 and 40.


Sensitivity Analysis Our calculations, which are based on the Freedman-Lane procedure under the assumption of partial linearity, rely on linear parametric approximations and on a particular choice of SES quantiles to define permutation orbits. Other choices are possible. Any or all of the four covariates that we use in the Freedman-Lane procedure under full linearity could have been used as conditioning variables to define restricted permutation orbits under a partial linearity assumption. We choose the SES index for nonparametric conditioning, since family background is known to be a powerful determinant of adult outcomes (see Cunha, Heckman, Lochner, and Masterov, 2006). Specifically, we use a dummy variable for

⁴⁹Statistics for female felony arrests are not shown in the table due to their low reliability: the small sample size and the low incidence of felony arrests.

⁵⁰The set of crime hypotheses is different for males and females due to small sample sizes: we cannot reliably measure the probability of incarceration for females for Perry sample.

⁵¹Heckman, Moon, Pinto, Savelyev, and Yavitz (2010) present a detailed analysis of total crime cost and its contributions to the economic return to the Perry program.







whether the SES index is above or below the sample median.

It is informative to conduct a sensitivity analysis on the effects of the choice of conditioning strata, which correspond to the covariates whose relationship with the outcome is assumed to be nonlinear rather than linear. To test the sensitivity of our results to the choice of stratum, we run a series of partially linear Freedman-Lane procedures with varying assumptions regarding the set of which covariates enter linearly.

The four pre-program covariates in question can be used either as Freedman-Lane regressors, which assume a linear relationship with outcomes, or as conditioning variables that limit the orbits of permutations to their selected quantiles, which allows for a nonlinear relationship. In Web Appendix F, we perform two types of sensitivity analyses. The first shows that the results reported in Tables 3–6 are robust to variations in the choice of SES index quantiles used to generate the strata on which permutations are restricted: median, tercile, or quartile. The second shows that our results are robust to the choice of which covariates enter the outcome model linearly.

Additional Evidence on the Effectiveness of the Perry Program In related work (Heckman, Moon, Pinto, Savelyev, and Yavitz, 2010), we calculate rates of return to determine the private and public returns to the Perry Preschool Program. We avoid the multiple hypothesis-testing problem by focusing on a single economically significant summary of the program. We use the conditioning approach adopted in this paper to control for compromised randomization. We find statistically significant rates of return for both males and females in the range of 6–10% per annum. This supports the evidence of substantial treatment effects presented in the current paper.

Understanding Treatment Effects While this paper tests for the existence of treatment effects due to the Perry Preschool Program, other recent work examines channels through which these beneficial effects are produced. Heckman, Malofeeva, Pinto, and Savelyev (2010) estimate a model of latent cognitive and noncognitive traits. In the early years during and

after the program, the IQ scores of treatment group participants surged, but by almost age 8, the treatment effect on IQ becomes nonexistent for males and relatively small for females. Their research shows that the effects of the Perry program arise primarily from boosts in noncognitive traits.

6 The Representativeness of the Perry Study

We next examine the representativeness of the Perry sample and characterize the target population within the overall African-American population. We construct a comparison group using the 1979 National Longitudinal Survey of Youth (NLSY79), a widely used, nationally representative longitudinal data set. The NLSY79 has panel data on wages, schooling, and employment for a cohort of young adults who were 14-22 at their first interview in 1979. This cohort has been followed ever since. For our purposes, an important feature is that the NLSY79 contains information on cognitive test scores as well as on noncognitive measures. It also contains rich information on family background. This survey is a particularly good choice for such a comparison as the birth years of its subjects (1957–1964) include those of the Perry sample (1957–1962). The NLSY79 also oversamples African Americans.

The Matching Procedure We use a matching procedure to create NLSY79 comparison groups for Perry control groups by simulating the application of the Perry eligibility criteria to the full NLSY79 sample. Specifically, we use the Perry eligibility criteria to construct samples in the NLSY79. Thus, the comparison group corresponds to the subset of NLSY79 participants who would likely be eligible for the Perry program if it were a nationwide intervention.

We do not have identical information on the NLSY79 respondents and the Perry entry cohorts, so we approximate a Perry-eligible NLSY79 comparison sample. In the absence of IQ scores in the NLSY79, we use Armed Forces Qualification Test (AFQT) scores as a proxy for IQ. We also construct a pseudo-SES index for each NLSY79 respondent using the available information.⁵²

We use two different subsets of the NLSY79 sample to draw inferences about the representativeness of the Perry sample. For an initial comparison group, we use the full African-American subsample in NLSY79. We then apply the approximate Perry eligibility criteria to create a second comparison group based on a restricted sub-sample of the NLSY79 data.

The U.S. population in 1960 was 180 million people, of which 10.6% (19 million) were African-American.⁵³ According to the NLSY79, the black cohort born in 1957–1964 is composed of 2.2 million males and 2.3 million females. We estimate that 17% of the male cohort and 15% of the female cohort would be eligible for the Perry program if it were applied nationwide. This translates into a population estimate of 712,000 persons out of the 4.5 million black cohort, who resemble the Perry population in terms of our measures of disadvantage.⁵⁴ For further information on the comparison groups and their construction, see Web Appendix H and Tables H.1 and H.2 for details.

How Representative is the Perry Sample of the Overall African-American Population of the United States? Compared to the unrestricted African-American NLSY79 subsample, Perry program participants are more disadvantaged in their family backgrounds. This is not surprising, given that the Perry program targeted disadvantaged children. Further, Perry participants experience less favorable outcomes later in life, including lower high school graduation rates, employment rates, and earnings. However, if we impose restrictions on the NLSY79 subsample that mimic the sample selection criteria of the Perry program, we obtain a roughly comparable group. Figure 5 demonstrates this comparability for parental highest grade completed at the time children are enrolled in the program. Web Appendix Figures H.1–H.5 report similar plots for other outcomes, including mother's age at birth,

⁵²For details, see the Web Appendix http://jenni.uchicago.edu/Perry/cost-benefit/reanalysis ⁵³See http://www.census.gov/population/www/documentation/twps0056/twps0056.html for more

details.

 $^{^{54}}$ When a subsample of the NLSY79 is formed using three criteria that characterize the Perry sample — low values of a proxy for the Perry socio-economic status (SES) index, low achievement test (AFQT) score, and non-firstborn status — this subsample represents 713,725 people in the United States. See Web Appendix H and Tables H.1 and H.2 for details.

earnings at age 27, and earnings at 40.⁵⁵ Tables H.1 and H.2 present additional details. The Perry sample is representative of disadvantaged African-American populations.

In Web Appendix I, we consider another aspect of the representativeness of the Perry experiment. Perry participants were caught up in the boom and bust of the Michigan auto industry and its effects on related industries. In the 1970s, as Perry participants entered the workforce, the male-friendly manufacturing sector was booming. Employees did not need high school diplomas to get good entry-level jobs in manufacturing, and men were much more likely to be employed in the manufacturing sector than women. The industry began to decline as Perry participants entered their late 20s.

This pattern may explain the gender patterns for treatment effects found in the Perry experiment. Neither treatments nor controls needed high school diplomas to get good jobs. As the manufacturing sector collapsed, neither group fared well. However, as noted in Web Appendix I, male treatment group members were somewhat more likely to adjust to economic adversity by migrating than were male controls, which may account for their greater economic success at age 40. The history of the Michigan economy helps to explain the age pattern of observed treatment effects for males, thereby diminishing the external validity of the study.

7 Relationship of This Study to Previous Research

Schweinhart et al. (2005) analyze the Perry data through age 40 using large sample statistical tests. They show substantial effects of the program for both males and females. They do not account for the compromised randomization of the experiment or the multiplicity of hypotheses tested. Heckman (2005) discusses the problems of the small sample size, the need to use small sample inference to analyze the Perry data, and the appropriate way to combine inference across hypotheses.

Anderson (2008) addresses the problem of multiple-hypothesis testing in the Perry data.

⁵⁵One exception to this pattern is that Perry treatment and control earnings are worse off than their matched sample counterparts.




Figure 5: Perry vs. NLSY79: Mean Parental Highest Grade Completed

Notes: Unrestricted NLSY79 is the full African-American subsample. Restricted NLSY79 is the African-American subsample limited to those satisfying the approximate Perry eligibility criteria: at least one elder sibling, Socio-economic status (SES) index at most 11, and 1979 AFQT score less than the African-American median. The reported "t" test is for the difference in means between the two populations.

He reanalyzes the Perry data (and data on other early childhood programs) using a stepdown multiple-hypothesis testing procedure due to Westfall and Young (1993). That procedure requires "subset pivotality," that is, that the multivariate distribution of any subvector of p-values is unaffected by the truth or falsity of hypotheses corresponding to p-values not included in the subvector. This is a strong condition.⁵⁶ Our method for testing multiple hypotheses is based on the stepdown procedure of Romano and Wolf (2005), which uses an assumption about monotonicity of the test statistics. Romano and Wolf (2005) show that their monotonicity assumption is weaker than the subset pivotality assumption.

Anderson applies permutation inference to avoid relying on asymptotically justified test statistics. We confirm his finding that even in the small Perry sample, asymptotic statistics are valid, so concerns about the use of large-sample inference to analyze the Perry samples are misplaced. However, in constructing his tests, Anderson assumes that a simple randomization was conducted in the Perry experiment. He does not address the problem of compromised randomization, neither does he correct for covariate imbalances between treatments and controls.

Anderson reports no statistically significant effects of the Perry program for males. We find that the Perry program improved the status of both genders on a variety of measures. One explanation for the difference between Anderson's conclusions and ours about the effectiveness of the program for males is that we adjust for covariate imbalances and compromised randomization while Anderson does not. As displayed in Tables 5 and 6, these adjustments sharpen the inference for males and lead to more rejections of the null hypothesis.

Another explanation for the contrast between our conclusions is differences in the blocks of variables used as the basis for the stepdown multiple-hypothesis testing procedures. To reduce the dimensionality of the testing problem, Anderson creates linear indices of outcomes at three stages of the life cycle. The outcomes used to create each index are quite diverse

 $^{^{56}}$ In Web Appendix D.3, we present an example, due to Westfall and Young (1993), where the subset pivotality condition is satisfied for testing hypotheses about means of a normal model but not for testing hypotheses about correlations.

and group a variety of very different outcomes (e.g., crime, employment, education). It is difficult to interpret his indices. Moreover, the components of his indices change with age. We conduct inference for interpretable blocks of hypotheses defined at different stages of the life cycle that are based on comparable outcomes (crime as one block, employment as another block, etc.).

8 Summary and Conclusions

Most social experiments are compromised by practical difficulties in implementing the intended randomization protocol. They also have a variety of outcome measures. This paper develops and applies a methodology for analyzing experiments as implemented and for generating valid tests of multiple hypotheses.

We apply our methods to analyze data from the Perry Preschool experiment. Evidence from the HighScope Perry Preschool Program is widely cited to support early childhood interventions. The consequences of imperfect randomization for inference are neglected by previous analysts of these data. This paper shows how to account for compromised randomization to produce valid test statistics.

Proper analysis of the Perry experiment also requires application of methods for smallsample inference and accounting for the large numbers of outcomes of the study. It is important to avoid the danger of artificially lowering p-values by selecting statistically significant outcomes that are "cherry picked" from a larger set of unreported hypothesis tests that do not reject the null.

We propose and implement a combination of methods to simultaneously address these problems. We account for compromises in the randomization protocol by conditioning on background variables to control for the violations of the initial randomization protocol and imbalanced background variables. We use small-sample permutation methods and estimate family-wise error rates that account for the multiplicity of experimental outcomes. The methods developed and applied here have applications to social experiments with small samples when there is imbalance in covariates between treatments and controls, reassignment after randomization, and multiple hypotheses.

The pattern of treatment response by gender varies with age. Males exhibit statistically significant treatment effects for criminal activity, later life income, and employment (ages 27 and 40), whereas female treatment effects are strongest for education and early employment (ages 19 and 27). There is, however, a strong effect of the program on female crime at age 40. The general pattern is one of strong early results for females, with males catching up later in life.

Our analysis of the representativeness of this program shows that Perry study families are disadvantaged compared to the general African-American population. However, application of the Perry eligibility rules to the NLSY79 yields a substantial population of comparable individuals. Based on the NLSY79 data, we estimate that the program targetted about 16% of the African-American population born during 1957–1964, which includes the birth years of the Perry participants.

We present some suggestive evidence that the limited effect of the Perry program on the education of males was due to the peculiarities of the Michigan economy. High school degrees were not required to work in well-paying manufacturing jobs. Perry treatment males appear to have adjusted to the decline in manufacturing that occurred in Michigan better than the controls. This accounts for the statistically significant treatment effects in employment and earnings found for males at age 40.

Few social experiments perfectly implement planned treatment assignment protocols. A proper analysis of such experiments requires recognizing the sampling plan as implemented. Our analysis shows that properly accounting for experiments as implemented can produce sharper results than analyses that proceed as if an ideal experiment was implemented.⁵⁷

References

- Anderson, M. (2008, December). Multiple inference and gender differences in the effects of early intervention: A reevaluation of the Abecedarian, Perry Preschool and early training projects. Journal of the American Statistical Association 103(484), 1481–1495.
- Anderson, M. J. and P. Legendre (1999). An empirical comparison of permutation methods for tests of partial regression coefficients in a linear model. *Journal of Statistical Computation and Simulation 62*, 271–303.
- Anderson, M. J. and J. Robinson (2001, March). Permutation tests for linear models. The Australian and New Zealand Journal of Statistics 43(1), 75–88.
- Benjamini, Y. and Y. Hochberg (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Association, Series* B 57(1), 289–300.
- Benjamini, Y., A. M. Krieger, and D. Yekutieli (2006, September). Adaptive linear step-up procedures that control the false discovery rate. *Biometrika* 93(3), 491–507.
- Beran, R. (1988a, September). Balanced simultaneous confidence sets. Journal of the American Statistical Association 83(403), 679–686.

⁵⁷In related work, Heckman, Pinto, Shaikh, and Yavitz (2009) take a more conservative approach to the problem of compromised randomization using weaker assumptions. Their inference is based on a partially identified model in which the distribution of D conditional on X is not fully known because the rules assigning persons to treatment are not fully determined. Unmeasured variables that determine assignment are also assumed to affect outcomes. They estimate conservative bounds for inference on treatment effects that are consistent with the verbal descriptions of the criteria used for reassignment.

This paper is less conservative than theirs because it adopts stronger assumptions: conditional exchangeability of treatment assignments within coarse strata of pre-program observables. As expected, this less conservative approach produces sharper inferences, although the inferences from the two approaches are in surprisingly broad agreement.

- Beran, R. (1988b, September). Prepivoting test statistics: A bootstrap view of asymptotic refinements. *Journal of the American Statistical Association* 83(403), 687–697.
- Campbell, F. A. and C. T. Ramey (1994, April). Effects of early intervention on intellectual and academic achievement: A follow-up study of children from low-income families. *Child Development 65*(2), 684–698. Children and Poverty.
- Campbell, F. A., C. T. Ramey, E. Pungello, J. Sparling, and S. Miller-Johnson (2002). Early childhood education: Young adult outcomes from the abecedarian project. Applied Developmental Science 6(1), 42–57.
- Cunha, F., J. J. Heckman, L. J. Lochner, and D. V. Masterov (2006). Interpreting the evidence on life cycle skill formation. In E. A. Hanushek and F. Welch (Eds.), *Handbook* of the Economics of Education, Chapter 12, pp. 697–812. Amsterdam: North-Holland.
- Fisher, F. M. (1966). The Identification Problem in Econometrics. New York: McGraw-Hill.
- Freedman, D. and D. Lane (1983, October). A nonstochastic interpretation of reported significance levels. *Journal of Business and Economic Statistics* 1(4), 292–298.
- Good, P. I. (2000). Permutation Tests: A Practical Guide to Resampling Methods for Testing Hypotheses (2 ed.). Series in Statistics. New York: Springer Verlag.
- Hanushek, E. and A. A. Lindseth (2009). Schoolhouses, Courthouses, and Statehouses: Solving the Funding-Achievement Puzzle in America's Public Schools. Princeton, NJ: Princeton University Press.
- Hayes, A. (1996, June). Permutation test is not distribution-free: Testing h_0 : $\rho = 0$. Psychological Methods 1(2), 184–198.
- Heckman, J. J. (1992). Randomization and social policy evaluation. In C. Manski and I. Garfinkel (Eds.), *Evaluating Welfare and Training Programs*, pp. 201–230. Cambridge, MA: Harvard University Press.

- Heckman, J. J. (2005). Invited comments. In L. J. Schweinhart, J. Montie, Z. Xiang,
 W. S. Barnett, C. R. Belfield, and M. Nores (Eds.), *Lifetime Effects: The High/Scope Perry Preschool Study Through Age 40*, pp. 229–233. Ypsilanti, MI: High/Scope Press.
 Monographs of the High/Scope Educational Research Foundation, 14.
- Heckman, J. J. (2010). The principles underlying evaluation estimators with an application to matching. Annales d'Economie et de Statistiques. Forthcoming.
- Heckman, J. J., H. Ichimura, J. Smith, and P. E. Todd (1998, September). Characterizing selection bias using experimental data. *Econometrica* 66(5), 1017–1098.
- Heckman, J. J., R. J. LaLonde, and J. A. Smith (1999). The economics and econometrics of active labor market programs. In O. Ashenfelter and D. Card (Eds.), *Handbook of Labor Economics*, Volume 3A, Chapter 31, pp. 1865–2097. New York: North-Holland.
- Heckman, J. J., L. Malofeeva, R. Pinto, and P. A. Savelyev (2010). Understanding the mechanisms through which an influential early childhood program boosted adult outcomes. Unpublished manuscript, University of Chicago, Department of Economics.
- Heckman, J. J., S. H. Moon, R. Pinto, P. A. Savelyev, and A. Q. Yavitz (2010, February). The rate of return to the HighScope Perry Preschool Program. *Journal of Public Economics* 94 (1-2), 114–128.
- Heckman, J. J. and S. Navarro (2004, February). Using matching, instrumental variables, and control functions to estimate economic choice models. *Review of Economics and Statistics* 86(1), 30–57.
- Heckman, J. J., R. Pinto, A. M. Shaikh, and A. Yavitz (2009). Compromised randomization and uncertainty of treatment assignments in social experiments: The case of Perry Preschool Program. Unpublished manuscript, University of Chicago, Department of Economics.

- Heckman, J. J. and J. A. Smith (1995, Spring). Assessing the case for social experiments. Journal of Economic Perspectives 9(2), 85–110.
- Heckman, J. J. and E. J. Vytlacil (2007). Econometric evaluation of social programs, part II: Using the marginal treatment effect to organize alternative economic estimators to evaluate social programs and to forecast their effects in new environments. In J. Heckman and E. Leamer (Eds.), *Handbook of Econometrics*, Volume 6B, pp. 4875–5144. Amsterdam: Elsevier.
- Herrnstein, R. J. and C. A. Murray (1994). The Bell Curve: Intelligence and Class Structure in American Life. New York: Free Press.
- Hotz, V. J. (1992). Designing an evaluation of the Job Training Partnership Act. In C. Manski and I. Garfinkel (Eds.), *Evaluating Welfare and Training Programs*, pp. 76–114. Cambridge, MA: Harvard University Press.
- Kurz, M. and R. G. Spiegelman (1972). The Design of the Seattle and Denver Income Maintenance Experiments. Menlo Park, CA: Stanford Research Institute.
- Lehmann, E. L. and J. P. Romano (2005). *Testing Statistical Hypotheses* (Third ed.). New York: Springer Science and Business Media.
- Meghir, C. and L. Pistaferri (2004, January). Income variance dynamics and heterogeneity. *Econometrica* 72(1), 1–32.
- Micceri, T. (1989, January). The unicorn, the normal curve, and other improbable creatures. Psychological Bulletin 105(1), 156–166.
- Pesarin, F. and L. Salmaso (2010). Permutation Tests for Complex Data: Theory, Application and Software. Chichester, UK: John Wiley and Sons.
- Reynolds, A. J. and J. A. Temple (2008). Cost-effective early childhood development programs from preschool to third grade. *Annual Review of Clinical Psychology* 4(1), 109–139.

- Romano, J. P. and A. M. Shaikh (2004). On control of the false discovery proportion. Technical Report 2004-31, Department of Statistics, Stanford University.
- Romano, J. P. and A. M. Shaikh (2006, August). Stepup procedures for control of generalizations of the familywise error rate. *Annals of Statistics* 34(4), 1850–1873.
- Romano, J. P. and M. Wolf (2005, March). Exact and approximate stepdown methods for multiple hypothesis testing. *Journal of the American Statistical Association* 100(469), 94–108.
- Rosenbaum, P. R. and D. B. Rubin (1983, April). The central role of the propensity score in observational studies for causal effects. *Biometrika* 70(1), 41–55.
- Schweinhart, L. J., H. V. Barnes, and D. Weikart (1993). Significant Benefits: The High-Scope Perry Preschool Study Through Age 27. Ypsilanti, MI: High/Scope Press.
- Schweinhart, L. J., J. Montie, Z. Xiang, W. S. Barnett, C. R. Belfield, and M. Nores (2005). Lifetime Effects: The High/Scope Perry Preschool Study Through Age 40. Ypsilanti, MI: High/Scope Press.
- The Pew Center on the States (2009, March). The facts. Response to ABC News Segements on Pre-Kindergarten. Available online at: http://preknow.org/documents/the_facts.pdf. Last accessed March 24, 2009.
- Weikart, D. P., J. T. Bond, and J. T. McNeil (1978). The Ypsilanti Perry Preschool Project: Preschool Years and Longitudinal Results Through Fourth Grade. Ypsilanti, MI: Monographs of the High/Scope Educational Research Foundation.
- Westat (1981). Impact of 1977 earnings of new fy 1976 ceta enrollees in selected program activities. *Continuous Longitudinal Manpower Survey. Net Impact Report 80-20*(1).
- Westfall, P. H. and S. S. Young (1993). Resampling-Based Multiple Testing: Examples and Methods for p-Value Adjustment. John Wiley and Sons.

Zhao, Z. (2004). Using matching to estimate treatment effects: Data requirements, matching metrics, and monte carlo evidence. The Review of Economics and Statistics 86(1), 91–107.